

REMARKS/ARGUMENTS

Claims 1, 5-7, 12-17 and 19 are now pending in this application.

Claim 12 has been amended to correct a minor typographical error. No new matter has been added.

Rejections under 35 U.S.C. §102 and §103

The only outstanding rejections were made over PCT publication WO 99/15652 to Miyata et al.

The Examiner has pointed out that the rejection can be overcome by filing a certified translation of the priority document thereby establishing that applicants had possession of the invention as of March 19, 1999. Applicants have attached hereto a certified copy of the priority document thereby overcoming the rejection.

Applicants appreciate the Examiner's indication that the earlier rejections have been withdrawn. In view of such and in view of the attached certified translation the application is believed to be in condition for allowance and an indication of such is respectfully requested.

The Examiner will note that the attached certified translation includes a verification statement indicating that the translator is well acquainted with both English and Japanese languages and that the attached document is a true and correct translation.

Applicants point out that Examples 7 and 8 on pages 34 through 36 of the attached translation disclose a diagnostic method. The diagnostic method is a method of diagnosing mesangial cell proliferative neuropathy. Such a method is claimed within claims such as claims 1 and 5 now pending in the application.

The examples 7-8 on pages 34-35 also disclose the reagent as well as the sequences such as those referred to within claim 6. The Examples 7-8 on pages 34-35 of the application are the working Examples 9 and 10 on pages 35-37 of the present application.

The method as claimed in claim 1 and the reagent as claimed in claim 6 which includes the specific SEQ IDs is encompassed by independent claim 12 which is supported at the same places in the specification.

Lastly the Examiner will note that kits useful in such methods as claimed within claim 17 are referred to in the translation such as in the Abstract on page 65 at lines 8-10.


In view of the attached verified translation of the priority document it is applicants position that the application is in condition for allowance and an indication of such is respectfully requested.

USSN: 09/936,883

The Commissioner is hereby authorized to charge any underpayment of fees associated with this communication, including any necessary fees for extensions of time, or credit any overpayment to Deposit Account No. 50-0815, order number SHIM-012.

Respectfully submitted,
BOZICEVIC, FIELD & FRANCIS LLP

Date: 5/May/05

By: 
Karl Bozicevic
Registration No. 28,807

Attachment: Certified Translation

BOZICEVIC, FIELD & FRANCIS LLP
1900 University Avenue, Suite 200
East Palo Alto, California 94303
Telephone: (650) 327-3400
Facsimile: (650) 327-3231



VERIFICATION OF TRANSLATION

I, Mariko Ogura

of 6th Fl., Kantetsu Tsukuba-Science-City Bldg. 1-1-1, Oroshi-machi, Tsuchiura, Ibaraki,
JAPAN

declare as follows:

1. That I am well acquainted with both the English and Japanese languages, and
2. That the attached document is a true and correct translation made by me to the best of my knowledge and belief of:

- (a) Japanese Patent Application No. Hei 11-075305
Entitled: "A METHOD FOR DETECTING MEGSIN PROTEIN AND USE
THEREOF"
Filed on March 19, 1999

April 26, 2005
(Date)

Mariko Ogura

(Signature of Translator)
Mariko Ogura



【Name of Document】 APPLICATION FOR PATENT

【Identification Number】 F2-101

【Filing Date】 March 19, 1999

【Addressee】 Esq. Commissioner of the Patent Office

【International Patent Classification】 G01N 33/53

【Inventor】

【Address】 4-2-3-101 Higashinaruse, Isehara-shi, KANAGAWA

【Name】 Toshio Miyata

【Applicant】

【Identification Number】 597142387

【Name or Appellation】 Kiyoshi Kurokawa

【Applicant】

【Identification Number】 597142376

【Name or Appellation】 Toshio Miyata

【Applicant】

【Identification Number】 000238201

【Name or Appellation】 Fuso Pharmaceutical Industries, Ltd.

【Attorney】

【Identification Number】 100102978

【Patent Attorney】

【Name or Appellation】 Hatsushi Shimizu

【Nominated Attorney】

【Identification Number】 100108774

【Patent Attorney】

【Name or Appellation】 Kazunori Hashimoto

【Payment】

【Registration Number】 041092

【Amount】 21000

【List of Attached Documents】

【Name of Document】 Description 1

【Name of Document】 Drawings 1

【Name of Document】 Abstract 1

【Proof】 Necessary

[Document Name] Specification

[Title of the Invention] A METHOD FOR DETECTING MEGSIN PROTEIN AND
USE THEREOF

[Claims]

5 [Claim 1] A method for evaluating renal functions, which
comprises measuring the megsin protein in a biological specimen.

 [Claim 2] The method for evaluating renal functions of claim
1, which comprises measuring the megsin protein in a biological
specimen, and comparing it with the megsin protein amount in a normal
10 specimen.

 [Claim 3] The method for evaluating renal functions of claim
1, wherein the biological specimen is urine.

 [Claim 4] The method for evaluating renal functions of claim
1, wherein the megsin protein in the biological specimen is measured
15 by an antigen-antibody reaction using an anti-megsin protein
antibody.

 [Claim 5] The method for evaluating renal functions of claim
4, wherein the anti-megsin protein antibody is a monoclonal antibody.

 [Claim 6] A reagent for diagnosing renal functions, which
20 comprises the anti-megsin protein antibody.

 [Claim 7] The reagent for diagnosing renal functions of claim
6, wherein the anti-megsin protein antibody is a monoclonal antibody.

 [Claim 8] A granule for detecting megsin protein in a
biological specimen, wherein the granule comprises a solid granule
25 to the surface of which an anti-megsin protein antibody is bound.

 [Claim 9] The granule for detecting megsin protein of claim
8, wherein the solid granule is magnetic.

 [Claim 10] The granule for detecting megsin protein of claim
8, wherein the relative density of the solid granule is not smaller
30 than 1.

 [Claim 11] The granule for detecting megsin protein of claim
8, wherein the anti-megsin protein antibody is a monoclonal antibody.

 [Claim 12] A method for detecting megsin protein in a
biological specimen, comprising the following steps of:

35 i) contacting the granule for detecting megsin protein with the
biological specimen, said granule comprising a solid granule to the

surface of which an anti-megsin protein antibody is bound;

ii) contacting the second anti-megsin protein antibody bound to a marker molecule with said granule for detecting megsin protein to which the biological specimen was contacted; and

5 iii) detecting the marker molecule bound to the megsin protein through the second anti-megsin protein antibody.

[Claim 13] The method for detection of claim 12, wherein the first anti-megsin protein antibody and the second anti-megsin protein antibody are both monoclonal antibodies.

10 [Claim 14] The method for detection of claim 13, wherein the first anti-megsin protein antibody and the second anti-megsin protein antibody are antibodies having different recognition sites.

[Claim 15] The method for detection of claim 12, wherein the biological specimen is urine.

15 [Claim 16] The method for detection of claim 12, wherein the biological specimen is blood.

[Claim 17] A kit for detecting megsin proteins, which comprises the following elements:

20 a) magnetic solid granules to which anti-megsin protein antibodies can be bound,

b) anti-megsin protein antibodies, which are bound to said magnetic solid granules in advance, or can be bound to them indirectly, and
c) a magnet.

[Claim 18] The kit for detecting megsin proteins of claim 25 17, further comprising an anti-megsin protein antibody to which a marker molecule is bound.

[Detailed Description of the Invention]

[0001]

[Technical Field of Industrial Application]

30 The present invention relates to a method for evaluating renal function, which comprises measuring the megsin protein in biological specimens such as urine, blood, and so on. Furthermore, the present invention relates to reagents for diagnosing renal function comprising anti-megsin protein antibodies.

35 [0002]

[Prior Art]

Kidney transplantation is the only method to treat terminal renal failure where the renal functions of hemofiltration and detoxication fail completely. However, the supply system supporting renal transplantations in Japan is far from well structured. In addition, social cognition on the transplantation method itself is not advanced. Therefore, the present status is that there is no alternative but to rely on dialysis as a renal replacement treatment.

Today, the number of patients who undergo dialysis is estimated to be approximately 170,000. An average annual treatment cost of about ¥6 million is needed per patient, and it is considered to be one of the biggest reasons that constrain the health insurance system. Furthermore, treatment with dialysis restraints the patient for 4 to 6 hours a day, 2 to 3 days a week, which greatly hampers the social activities of the patient.

Renal failure is a condition patients with renal diseases finally reach. The cause and background leading to renal failure is diverse. There are many cases in which diseases that usually originate outside the kidneys such as drug poisoning, infectious diseases, malignant tumors, diabetes, systemic lupus erythematosus (SLE), cause renal disorders and finally lead to renal failure.

Significant subjective symptoms of renal disorders appear only at late stages, namely only after the condition has almost worsened to renal failure. Therefore, they are easily overlooked, and numerous cases exist where at the point when symptoms appear, the kidney has already suffered an unrecoverable damage. Thus, it is important to discover renal disorders at an earlier stage as possible before the expression of subjective symptoms in order to prevent transition to renal failure, and also to eliminate constraints dialysis treatment puts on public medical insurance finances.

[0003]

Up to now, the examination of urinary proteins and urinary deposits, the so-called urinalysis, was widely carried out to diagnose renal disorders. However, urinary proteins increase temporarily

even in normal healthy persons due to extreme exercise, psychological stress, abundant meat diet, before menstruation, and so on. On the other hand, there are urinary proteins that are unrelated to renal diseases, such as orthostatic albuminuria found mainly in adolescents
5 (in around 0.5% of normal healthy persons). Urinary proteins are also detected in uropathies, bladder disorders, female genital tract disorders, and so on. Thus, it is difficult to make a definite diagnosis of renal disorders only by examining urinal proteins.

Urine deposits are observed with a microscope after
10 centrifugation of urine. However, erythrocyte deposits are also observed in the urine deposits of normal healthy persons, and may also derive from disorders other than renal disorders, such as those of organs related to the urinary tract. Therefore, urine deposits are also not sufficient enough to make definite diagnoses of renal
15 disorders.

A method to diagnose diabetic nephropathy at an early stage is known, which comprises the following steps: (1) assaying albumin that leaked into the urine as an indicator of diabetic nephropathy; and (2) comparing the value with the normal value of a normal healthy
20 person. However, the exact state of diabetic nephropathy cannot be understood since the albumin content in the urine changes even in a normal healthy person.

In addition, serum creatinine (Cr) and blood urea nitrogen (BUN) are assayed to examine the retention of urinary components
25 in the blood, but this assay is also easily influenced by diet. Thus, abnormal values in assays of urinary proteins, as well as serum Cr and BUN does not necessarily imply a renal disorder, and they may frequently occur in normal healthy persons or patients with other diseases.

30 [0004]

Further, diagnosis of renal disorders by measuring various substances including urinary β_2 -microglobulin, N-acetylglucosaminidase (NAG), IgG, transferrin, interleukin-6, and such are being tried. However, there are many cases where these
35 measurements do not correspond to the severity of the renal disorder, and therefore, are far from being effective.

Moreover, although a disorder of the whole kidney or involvement of immune reactions may be guessed by measuring these blood components in the urine, it is difficult to identify the site within renal tissue that is affected by the disorder. No examination method other than those mentioned above is known for diagnosing renal disorders and determining the severity with a sufficient sensitivity and specificity. Histological diagnosis by renal biopsy is regarded essential for ultimately diagnosing and determining the severity of renal disorders.

[0005]

However, renal biopsy is an invasive examination, and frequently accompanies dangers of complications such as hemorrhage, infections, and so on. Further, to conduct the examination, hospitalization in a well-equipped facility having specialists is required, and the physical and social burden on the patient cannot be ignored.

As mentioned above, the examination by urinalysis is simple and convenient, and is also an excellent examination method that enables treatment of a large amount of specimens. However, from the perspective of providing a definite diagnosis of renal disorders, it is far from being satisfactory. On the other hand, renal biopsy is an authentic method to diagnose and determine the severity of renal disorders. However, its use is highly restricted, which cannot be helped. On this account, a method to diagnose renal disorder, which has the simplicity and convenience of a urinalysis, as well as the accuracy of a renal biopsy, has been desired.

[0006]

Proteins specifically expressed in specific tissues, not only those in the kidney, are often used as indicators of functional disorders of the organ. For example, enzymatic proteins such as LDH and γ GTP are widely used as markers of hepatic function. However, no protein specific to the kidneys is known which serves as an indicator of their functions.

The present inventors isolated a gene called megsin as a gene especially strongly expressed in mesangial cells by a macroscale DNA sequence analysis and database analysis, and succeeded in obtaining

the megsin protein comprising 380 amino acids encoded by the megsin full-length cDNA clone. The present inventors, further found that the human megsin protein belongs to the SERPIN (serine protease inhibitor) superfamily (R.Carrell et al., Trends Biochem.Sci., 10:20 (1985); R.Carrell et al., Cold Spring Harbor Symp.Quant.Biol., 52:527 (1987); E.K.O.Kruithof et al., Blood, 86:4007 (1995); J.Potempa et al., J.Biol.Chem., 269:15957 (1994); E.Remold-O'Donnell, FEBS Let., 315:105 (1993)) according to an amino acid homology search by the FASTA program using the Swiss Prot database (T.Miyata et al. J.Clin.Invest., 120:828-836 (1998)). These findings were filed as a patent application (PCT/JP98/04269).

Human megsin protein expression was weak in human fibroblasts, smooth muscle cells, endothelial cells, and keratinocytes, and was strong especially in mesangial cells. The comparison of megsin protein expression level in renal tissues from IgA nephropathy patients and normal healthy individuals revealed that the expression level of megsin protein in IgA nephropathy patients was significantly higher. Accordingly, it is possible that megsin protein existing in the urine or blood is useful as a marker of mesangial cell proliferative nephropathy, for example, nephropathies such as IgA nephropathy. There is a possibility that megsin protein gene expression is deeply related with the onset and progression of renal diseases. However, the correlation between megsin protein gene expression and progression of renal disease state was unknown.

[0007]

[Problems to Be Solved by the Invention]

An objective of the present invention is to solve problems such as those above, and to provide a method that enables diagnosis of renal diseases and determination of their severity, and also to provide reagents and reagent kits used for the method.

[0008]

[Means to Solve the Problems]

Firstly, the present inventors thought that it is necessary to measure specific proteins related to the state of the disease to make a definite diagnosis of renal disorders and determine their severity, and focused their attention on mesangial cells, which exist in the

glomerulus. The mesangium is mapped on the lobule center in the glomerular tuft and is a tissue that functions as a core to bind the lobules together. The mesangium is covered by the glomerular basement membrane. It is composed of cells (mesangial cells), which are divided from the capillary lumen by endothelial cells, and a three-layered amorphous material (mesangial matrix), which is in sequence with the lamina rara interna in the glomerular basement membrane.

[0009]

The mesangial cells play a pivotal role in maintaining the structure and function of the glomerulus. Their proliferation is thought to be a main factor in the onset of glomerular disorders such as glomerular nephritis and glomerulosclerosis. Furthermore, mesangial cells are targets damaged by all types of nephritis. For example, proliferation of mesangial cells and accumulation of extracellular mesangial matrix is thought to be the first step in the development of glomerulosclerosis in patients suffering from various glomerular diseases, such as chronic nephritis and diabetic nephritis, the two main causes of end-stage renal failure(D. Schlondorff, *Kidney Int.*, 49: 1583-1585, (1996); R. B. Sterzel et al., *Glomerular mesangial cell. Immunologic Renal Diseases*, 595-626 (1997)) Therefore, it is thought that identification of a gene expressed specifically in mesangial cells and elucidation of its function would be helpful in understanding the biological characteristics of mesangial cells and the causes of diseases relating to mesangial cells, and in turn, treating or diagnosing diseases related to mesangial cells.

It has been reported that steroidal agents are effective for IgA nephropathy with an extreme proliferation of mesangial cells (J.V.Donadio et al., *J.Am.Soc.Nephrol.* 8:1324-1332 (1997)). However, use of steroidal agents often causes serious side effects, and thus, aimless administration of steroidal agents to types of IgA nephropathy for which steroidal agents are ineffective, should be avoided. Therefore, a matching up with the severity of the tissue such as the degree of proliferation of mesangial cells, and so on, is essential for steroid therapy. Accordingly, a renal function assay that can

simply and conveniently evaluate the degree of proliferation of mesangial cells was long desired.

The present inventors assumed that megsin protein gene expression increases in relation to the onset and progress of renal diseases, and in association thereto, megsin protein production increases and leads to a leakage of megsin protein into urine or blood, the amount of which may be related to the progression of disease state. To verify the mechanism, the inventors attempted to measure and compare the concentration or amount of megsin protein in various biological specimens, and thus, accomplished the present invention through the result that renal diseases related to megsin protein can be evaluated by measuring the concentration or amount of the megsin protein. Thus, the present invention relates to a method for evaluating renal functions, as well as to reagents or reagent kits for the method, which are as follows:

- (1) a method for evaluating renal functions, which comprises measuring the megsin protein in a biological specimen;
- (2) the method for evaluating renal functions of (1), which comprises measuring the megsin protein in a biological specimen, and comparing it with the megsin protein amount in a normal specimen;
- (3) the method for evaluating renal functions of (1), wherein the biological specimen is urine;
- (4) the method for evaluating renal functions of (1), wherein the megsin protein in the biological specimen is measured by an antigen-antibody reaction using an anti-megsin protein antibody;
- (5) the method for evaluating renal functions of (4), wherein the anti-megsin protein antibody is a monoclonal antibody;
- (6) a reagent for diagnosing renal functions, which comprises the anti-megsin protein antibody;
- (7) the reagent for diagnosing renal functions of (6), wherein the anti-megsin protein antibody is a monoclonal antibody;
- (8) a granule for detecting megsin protein in a biological specimen, wherein the granule comprises a solid granule to the surface of which an anti-megsin protein antibody is bound;
- (9) the granule for detecting megsin protein of (8), wherein the solid granule is magnetic;

(10) the granule for detecting megsin protein of (8), wherein the relative density of the solid granule is not smaller than 1;

(11) the granule for detecting megsin protein of (8), wherein the anti-megsin protein antibody is a monoclonal antibody;

5 (12) a method for detecting megsin protein in a biological specimen, comprising the following steps of:

i) contacting the granule for detecting megsin protein with the biological specimen, said granule comprising a solid granule to the surface of which an anti-megsin protein antibody is bound;

10 ii) contacting the second anti-megsin protein antibody bound to a marker molecule with said granule for detecting megsin protein to which the biological specimen was contacted; and,

iii) detecting the marker molecule bound to the megsin protein through the second anti-megsin protein antibody;

15 (13) the method for detection of (12), wherein the first anti-megsin protein antibody and the second anti-megsin protein antibody are both monoclonal antibodies;

(14) the method for detection of (13), wherein the first anti-megsin protein antibody and the second anti-megsin protein antibody are
20 antibodies having different recognition sites;

(15) the method for detection of (12), wherein the biological specimen is urine;

(16) the method for detection of (12), wherein the biological specimen is blood;

25 (17) a kit for detecting megsin proteins, which comprises the following elements:

a) magnetic solid granules to which anti-megsin protein antibodies can be bound,

30 b) anti-megsin protein antibodies, which are bound to said magnetic solid granules in advance, or can be bound to them indirectly, and

c) a magnet;

(18) the kit for detecting megsin proteins of (17), further comprising an anti-megsin protein antibody to which a marker molecule is bound.

[0010]

35 [Mode for Carrying Out the Invention]

As mentioned above, the megsin protein was isolated as a protein

encoded by a gene that is highly expressed in human kidney mesangial cells. The megsin protein of the present invention includes not only a protein having the amino acid sequence of SEQ ID NO: 2 (human megsin protein), but also proteins functionally equivalent thereto. For example, megsin protein homologues from species other than human can be mentioned as proteins that are functionally equivalent. Other examples are precursors and fragments of these homologues. Megsin protein-protease complexes can also be used as indicators of renal functions, as long as they can be detected within biological specimens. Megsin protein homologues of species other than humans can be used as indicators for the evaluation of renal functions in the respective species. Thus, they are useful to understand disease states in renal function disorder models.

[0011]

Megsin protein homologues from species other than human can be obtained, for example, as follows: First, the homologue cDNA is amplified by PCR utilizing a degenerative primer designed based on the nucleotide sequence of the cDNA encoding the human megsin protein of SEQ ID NO: 1 to correspond to the beginning and end of the ORF. In case where the amplified cDNA is not a full-length cDNA, its upstream nucleotide sequence can be determined by methods such as 5'RACE. For example, RT-PCR is conducted using the mRNA, which is extracted from cultured mesangial cells, as the template for rats. Degenerative primers used can be, for example, the oligonucleotides with the following nucleotide sequences. Part of the structure of the rat homologue megsin protein can be revealed by sequence determination of the clone obtained by such a PCR method.

Degenerative primer FY: GTGAATGCTGTGTACTTAAAGGCAANTGN/ SEQ ID NO: 3 (corresponding to 172VNAVYFKGK180)

Degenerative primer R21: AANAGRAANGGRTCNGC/ SEQ ID NO:4 (wherein R is A or G; corresponding to 357ADHPFLF363)

However, to obtain the 5' regions, it is necessary to conduct a degenerate PCR again by preparing gene specific primers from clone fragments of the rat megsin protein, since only a part of the cDNA can be amplified with these primer pairs. The nucleotide sequences of the primers used, which correspond to the N terminal amino acid

sequences of the megsin proteins are as follows:

Degenerative primer RM-CtermC1: ATGGCNTCNGCNGCNGCNGCNAAYGC/ SEQ ID NO:5 (wherein Y is T or C; corresponding to the sequence encoding the N-terminus of the megsin protein)

5 RM-MR-A2: CGACCTCCAGAGGCAATTCCAGAGAGATCAGCCCTGG/ SEQ ID NO: 6

RM-MR-A1: GTCTTCCAAGCCTACAGATTTCAAGTGGCTCCTC/ SEQ ID NO: 7

(Each of them is a rat megsin protein specific reverse primer)

The 3' region of the cDNA encoding the rat homologue can be derived by inserting the obtained PCR product into the pGEM-T-easy
10 vector (Promega) to determine the nucleotide sequence. Finally, the nucleotide sequence of the full-length cDNA encoding the homologue can be determined by conducting 5'-RACE and 3'-RACE using a primer designed based on the nucleotide sequence obtained as above. Two gene specific antisense primers, namely RM-PRO1:
15 GCTCAGGGCAGTGAAGATGCTCAGGGAAGA/ SEQ ID NO:8 and RM-PRO2: CTGACGTGCACAGTCACCTCGAGCACC/ SEQ ID NO:9, can be used for the 5'-RACE for rats. On the other hand, gene specific sense primer RM-MR-S3: GAGGTCTCAGAAGAAGGCACTGAGGCAACTGCTGCC/ SEQ ID NO:10 can be used in the 3'-RACE. Mouse homologues can be isolated by the same method.
20 The sequence of rat and mouse megsin cDNA nucleotide sequence depicted in SEQ ID NO:15 (rat) and SEQ ID NO:17 (mouse) were determined according to the method above.

[0012]

The method used in the present invention to measure the megsin
25 protein is not restricted. For example, immunological assays, which make use of the immunological reaction between an antibody for the megsin protein and the megsin protein, are excellent in both specificity and sensibility. Furthermore, the megsin protein can be measured by the Western Blot method utilizing an antibody against
30 the megsin protein. Since the megsin protein is a protease inhibitor, it can be detected using its inhibition activity on protease as the indicator. Alternatively, the megsin protein can be assayed utilizing its affinity towards the protease.

[0013]

35 The source or preparation method of the antibody, which is essential for immunological assays of the megsin protein, can be any

so long as the antibody can recognize the detection target, which is the megsin protein. Thus, antibodies to be used include polyclonal antibodies, monoclonal antibodies, mixtures thereof, etc. Furthermore, fragments containing the variable region of the antibody molecule are included. The antibody against megsin protein can be obtained, for example, as follows. For example, antibodies used in the present invention comprise those antibodies raised against the protein having the amino acid sequence of SEQ ID NO:2. The antibody (for example, polyclonal antibody or monoclonal antibody) or the antiserum against the megsin protein or partial amino acids sequence thereof can be prepared according to conventional methods for antibody or antiserum preparation using megsin proteins, oligopeptides comprising a partial amino acid sequence thereof, as well as fusion proteins, such as c-myc-(His)₆-Tag-megsin protein and MBP-megsin protein, as the antigens. For example, monoclonal antibodies can be prepared following the methods below. In cases where synthetic peptides with partial amino acid sequence are used as immunogens, it is generally convenient to use highly hydrophilic amino acid sequences that are specific to megsin protein. Such amino acid sequences include those shown in SEQ ID NO:11 to 14, which were used in the examples.

[0014]

The megsin protein of the present invention or a synthetic peptide having a partial amino acid sequence of the present megsin protein, is administered alone or together with a carrier or diluent to a warm-blooded animal at a site capable of producing antibodies. Synthetic peptides used as immunogens are in the form peptides bound to carrier proteins such as bovine thyroglobulin and keyhole limpet hemocyanin. To enhance the antibody productivity, the complete Freund's adjuvant or incomplete Freund's adjuvant can be administered together with the antigen. Immunization is performed once every 1 to 6 weeks, a total of about 2 to 10 times, in general. Warm-blooded animals to be used are, for example, monkeys, rabbits, dogs, guinea pigs, mice, rats, sheep, goats, and domestic fowls, preferably mice or rats. Monoclonal antibody-producing cells can be prepared by selecting immunized warm-blooded animals, such as mice, in which an

antibody titer is detected, obtaining the spleen or lymph nodes from the animals 2 to 5 days after the final immunization, and fusing the antibody producing cells contained in these tissues with myeloma cells. Reacting the labeled megsin protein described below with the antiserum, and measuring the activity of the label bound to the antibody enables measurement of the antibody titer in the antiserum.

[0015]

A monoclonal antibody of the present invention that does not crossreact with proteins other than megsin protein can be obtained by selecting an antibody that recognizes epitopes specific to megsin protein. In general, an epitope specific to the protein comprises at least 7 or more continuous amino acid residues, preferably 10 to 20 amino acids in the amino acid sequence of the protein. Therefore, for example, a monoclonal antibody recognizing an epitope comprising peptides having an amino acid sequence selected from the amino acid sequence of SEQ ID NO: 2, SEQ ID NO: 16(rat), or SEQ ID NO: 18(mouse), and comprising at least 7 continuous amino acid residues can be a monoclonal antibody specific to the megsin protein. Conserved amino acid sequences among the amino acid sequences of SEQ ID NO: 2, SEQ ID NO: 16(rat), and SEQ ID NO: 18(mouse) can be selected to choose epitopes common to the megsin protein family. If a region contains amino acid sequences specific to all sequences, a monoclonal antibody capable of recognizing different species can be selected.

An anti-MEGSIN protein monoclonal antibody can be separated and purified according to a separation and purification method for immunoglobulin, similar to the separation and purification of polyclonal antibodies. The known purification methods include, for example, salting out, alcohol precipitation, isoelectric point precipitation, electrophoresis, adsorption and desorption method by ion exchangers (for example, DEAE), ultra centrifugation, gel filtration, or specific purification methods whereby an antibody is exclusively collected, for example, methods whereby an antigen is collected using an antigen binding solid phase or active adsorbent, such as Protein A or Protein G, and thereafter the binding is dissociated to obtain the antibody.

[0016]

Monoclonal antibodies and polyclonal antibodies obtained in such a manner that recognize the megsin protein of the present invention, can be used for the diagnosis and treatment of diseases relating to mesangial cells. Examples of methods that use these
5 antibodies to measure the megsin protein include a sandwich assay, which comprises reacting the megsin protein with an antibody binding to an insoluble carrier and a labeled antibody and detecting MEGSIN protein in the sandwiched complex produced by the reaction, or a competitive method comprising competitively reacting labeled human
10 urine-derived megsin protein and human urine-derived megsin protein in a sample with an antibody to measure human urine-derived megsin protein in the sample based on the labeled antigen amount that reacted with the antibody.

The assaying of human urine-derived megsin protein by the
15 sandwich method is conducted by a 2 step method in which an immobilized antibody is reacted with human urine-derived megsin protein, unreacted materials are completely removed by washing, and a labeled antibody is added to form the immobilized antibody-labeled human urine-derived MEGSIN protein antibody complex, or one step
20 method in which the immobilized antibody, the labeled antibody, and human urine-derived MEGSIN are mixed at the same time.

[0017]

Examples of insoluble carriers used for the assay include, for example, polystyrene, polyethylene, polypropylene, polyvinyl
25 chloride, polyester, polyacrylate, nylon, polyacetal, synthetic resin such as fluoride resin, etc., polysaccharides such as cellulose, agarose, etc., glass, metals, etc. The form of the insoluble carrier can be varied and includes a shape like a particle, tray, sphere, fiber, stick, board, container, cell, test tube, etc.
30 The antibody-adsorbed carrier should be stored in a cool place in the presence of appropriate preservatives, such as sodium azide.

On the other hand, solid granules are known as reagents used to evaluate the fertility of human sperm cells. These granules comprise on their surfaces monoclonal antibodies that specifically
35 react with human sperm cells having undergone the acrosome reaction (Japanese Patent No.2651249). This granule used for sperm fertility detection is a solid granule to which a monoclonal antibody that

specifically reacts with human sperm cells having undergone the acrosome reaction is bound. The fertility of human sperm cells can be evaluated by reacting sperm cells with the granules and counting sperm cells bound to the granules. The fertility can be tested readily and safely without the use of complicated techniques like radioactivity, fluorescence, and so on. Samples of small quantities can be assayed by using magnetic solid granules since the granules can be aggregated easily by using a magnet. The present inventors found out that it is possible to detect megasin protein more easily and accurately by applying this technique.

[0018]

Antibodies can be immobilized by known chemical binding or physical adsorption methods. Chemical binding methods include, for example, a method using glutaraldehyde, the maleimide method using N-succinimidyl-4-(N-maleimidomethyl)cyclohexane-1-carboxylate, N-succinimidyl-2-maleimidoacetate, etc., and the carbodiimide method using 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride, etc. Further examples are, the maleimidobenzoyl-N-hydroxysuccinimidoester method, the N-succinimidyl-3-(2-pyridyldithio)propionate method, the bisdiazolated benzidine method, and dipalmityllysine method. Alternatively, a complex, which is prepared by reacting two different antibodies having different epitopes with the substance to be detected, can be captured with a third antibody immobilized by the above method.

[0019]

The label is not limited, so long as it is useful in immunoassays. Specifically, enzymes, fluorescent substances, luminescent substances, radioactive substances, metal chelates, etc. can be used. Preferable labeling enzymes are, for example, peroxidase, alkaline phosphatase, β -D-galactosidase, malate dehydrogenase, *Staphylococcus* nuclease, delta-5-steroid isomerase, α -glycerol phosphate dehydrogenase, triosephosphate isomerase, horseradish peroxidase, asparaginase, glucose oxidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase, and acetylcholine esterase, etc. Preferable

fluorescent substances include, for example, fluorescein isothiocyanate, phycobiliprotein, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, and orthophthalaldehyde. Preferable luminescent substances include, for example, isoluminol, lucigenin, luminol, aromatic acridinium ester, imidazole, acridinium salt and its modified ester, luciferin, luciferase, and aequorine. Preferable radioactive substances include, for example, ^{125}I , ^{127}I , ^{131}I , ^{14}C , ^3H , ^{32}P , ^{35}S , etc.

[0020]

Methods for binding the above labels are known. Specifically, direct and indirect labeling can be used. The commonly used direct labeling method is the method in which an antibody or an antibody fragment is chemically covalent-bound to a label using a cross-linking agent. Cross-linking agents include N,N'-orthophenylenedimaleimide, 4-(N-maleimidomethyl)cyclohexanoate N-succinimide ester, 6-maleimidohexanoate N-succinimide ester, 4,4'-dithiopyridine, and other known crosslinking agents. The crosslinking agent can be reacted with enzymes and antibodies by known methods depending on the characteristics of the cross-linking agent. An example of the indirect labeling method comprises binding an antibody to a low molecular weight hapten such as biotin, dinitrophenyl, pyridoxal, or fluorescamine, and indirectly labeling the antibody with the binding partner to the hapten. Avidin and streptavidin can be used as a recognition ligand for biotin, whereas dinitrophenyl, pyridoxal, or fluorescamine are labeled with antibodies recognizing these haptens.

Horseradish peroxidase can be used as an enzyme for labeling antibodies. This enzyme is useful because it can react with many substrates and be easily bound to antibodies by the periodate method. Occasionally, antibody fragments, for example, Fab', Fab, F(ab')₂ are used as antibodies. Both polyclonal and monoclonal antibodies can be labeled with an enzyme by the same method. Enzyme-labeled antibodies obtained using the above crosslinking agent can be purified by known methods such as affinity chromatography, etc. to serve as more sensitive immunoassay systems. Purified enzyme-labeled antibodies are stored with a preservative such as

thimerosal and a stabilizer such as glycerol. Labeled antibodies can be lyophilized and stored in the cool and dark place for a long time.

[0021]

5 When the label is an enzyme, its substrate and, if necessary, a coloring agent is used for measuring its activity. When peroxidase is used as an enzyme, H_2O_2 is used as the substrate solution, and 2,2'-azino-di-[3-ethylbenzothiazolinesulfonic acid] ammonium salt (ABTS), 5-aminosalicylic acid, 10 orthtophenylenediamine, 4-aminoantipyrine, or 3,3', 5,5'-tetramethylbenzidine, etc. are used as coloring agents. When alkaline phosphatase is used as the enzyme, orthonitrophenylphosphate, paranitrophenylphosphate, etc. can be used as substrates. When β -D-galactosidase is used as the enzyme, 15 fluorescein-di-(β -D-galactopyranoside), 4-methylumbelliferyl- β -D- galactopyranoside, etc. can be used as substrates. The present invention also includes an immunoassay reagent for megsin comprising labeled or immobilized monoclonal or polyclonal antibodies. The present invention further includes a kit 20 comprising this reagent and an indicator for detection labels and a control sample, etc.

Any biological sample such as body fluid including blood plasma, serum, blood, urine, tissue fluid, or cerebrospinal fluid, etc. can be used as samples for measuring the megsin protein of the 25 present invention, as long as they contain megsin protein or its precursor or a fragment. Among these biological samples, megsin protein can be detected with high frequency especially in urine, accompanied by a proliferation and activation of mesangial cells. Therefore, measurement of megsin in urine is useful as a marker for 30 mesangial proliferative nephropathy, such as IgA nephropathy.

[0022]

Evaluation of renal functions as used herein means to comprehend the condition of mesangial cells, which is an important cell constructing kidney tissue, and to determine the presence or severity 35 of a renal disease causing abnormalities in mesangial cells. Renal disease as used herein refers to any disease causing an abnormality

in mesangial cells. More specifically, IgA nephropathy, acute glomerulonephritis, focal sclerosing glomerulopathy, membranoproliferative glomerulonephritis, diabetic nephritis, lupus nephritis, and so on, can be mentioned as such diseases. The condition of renal functions affected by these renal diseases can be evaluated by the method of the present invention. Among the above diseases, the present method for evaluating renal function is especially useful as a marker for mesangial cell proliferative nephropathy, such as the IgA nephropathy. Moreover, the method of the present invention for evaluating renal function may be applied not only to determine the existence or severity of a renal disease, but also to evaluate a therapeutic effect or make a prognosis. In addition, the method of the present invention for evaluating renal function enables the exclusion of diseases not ascribable to the proliferation of mesangial cells like acute pyelonephritis, chronic pyelonephritis, minimal-change nephritic syndrome, chronic glomerulonephritis, renal amyloidosis, etc.

[0023]

To evaluate renal functions according to the present invention, a biological specimen from an individual whose renal function should be tested is obtained and the concentration of megsin protein therein is measured following the above-mentioned methods. Preferably, the amount of megsin protein is determined from the measured concentration and the volume, and compared to that of a normal healthy person. To calculate the amount of megsin protein using urine as the sample, all urine of a day is pooled and the amount of urine is measured. Thereby, it is possible to determine the amount of megsin protein in a day's urine. Alternatively, it is possible to presume the amount by creatinine correction, even when the urine is sampled occasionally. The term "creatinine correction" refers to a method to correct the influence of dilution (or concentration) of the object component due to the changes in urine amount based on the creatinine concentration. The creatinine amount excreted daily into the urine is constant. Based on this fact, the percentage occasionally collected urine occupies in the daily total excretion is calculated from the creatinine concentration, and the concentration of the object

component in the same urine is converted into the total excretion of a day. Alternatively, correction methods generally used in the diagnosis of renal functions such as weight correction is applied to presume the amount in blood. Weight correction is a method, wherein
5 the amount of blood component is calculated based on the volume presumed from the weight of the individual from whom the blood was taken.

Alternatively, changes in renal function can be also followed in a time course without having to convert into amounts by observing
10 the changes of megsin protein concentration in a biological specimen derived from a certain individual. Alternatively, an abnormality in renal function can be determined by configuring the normal values for megsin protein concentration in body fluids of specific species or groups like the race, and then, comparing the megsin protein
15 concentration (or amount) of an individual with the normal value.

The biological specimen used in the present invention can be urine or blood. Urine is more preferable since it can be collected in a noninvasive manner and is at the same time a sample that reflects the condition of renal functions directly. Although blood must be
20 collected with some invasiveness, since megsin protein is a protein specific to the kidney, an abnormality of the measured value closely relates to an abnormality of renal function. Thus, a high specificity can be expected if the amount of megsin protein in blood and urine is used as indicators of renal function. Compared to the currently
25 widely used method of screening for renal function disorder using urinary proteins as indicators, a more specific screening method with a superior sensibility can be expected by utilizing the megsin protein as the indicator. A method for evaluating renal functions by measuring the megsin protein in urine based on the immunological
30 method for measuring the megsin protein using a monoclonal antibody will be explained in detail below.

[0024]

(A) Preparation of the antibody

(1) Immunization of the animal and preparation of antibody-producing
35 cells

The immunization of the animal is conducted, for example, as

follows. A mammal, such as a rat, mouse, and so on, is immunized with the human megsin protein purified according to conventional methods (for example, T.Miyata et al., J.Clin.Invest., 120:828-836 (1998)). It is preferable to immunize animals of the same strain as the animal strain from which the permanently proliferating cell used during cell fusion is obtained. For example, mice at the age of 8 to 10 weeks are preferable. Mice of both genders will do. The immunization is conducted as follows: (1) the purified human megsin protein is mixed to an emulsion with an appropriate adjuvant (for example, Freund's complete adjuvant, aluminum hydroxide gel-pertussis vaccine, etc.), and (2) administered to the animal subcutaneously, intraperitoneally, intravenously, etc. Repeating this immunization process 2 to 5 times, with an interval of 1 to 2 weeks. The final immunization is accomplished by an intraperitoneal administration of 0.5 to 2 μ g human megsin protein to the animal. Polyclonal antibodies are obtained from the body fluid of the thus immunized animal. 3 to 7 days after each immunization process, blood is collected from venus plexus of the eyeground, and the antibody titer of the serum is measured by the protein A rosette assay (Eur.J.Immunol., 4:500-507 (1974)) described below. The antibody or the antibody-producing cell is harvested as the antibody titer had elevated enough.

[0025]

The protein A rosette assay is, for example, performed as follows: (1) 72 well Terasaki Plate (Falcon) is coated with the human erythroblast cell line K562 (Japanese Cancer Research Resources Bank (JCRS)); (2) samples diluted with PBS (sodium dihydrogen phosphate 2.90 g, potassium monohydrogen phosphate 0.20 g, sodium chloride 8 g, potassium chloride 0.2 g, distilled water 1 L) are added; (3) the plate is left standing for 30 minutes at 37°C in a CO₂ incubator; (4) the wells are washed with PBS; (5) sheep erythrocytes coated with protein A (Amersham Pharmacia Biotech) are added; and, (6) the formed rosettes are observed with a microscope.

Antibody-producing cells are harvested from the above human megsin protein immunized animal. Antibody-producing cells can be obtained from the spleen, lymph nodes, peripheral blood, and so on. However, the spleen is especially preferred. For example, 3 to 4 days

after the final immunization, the spleen is excised aseptically, cut in a Minimal Essential Medium (MEM) (Nissui Pharmaceutical), detach tissues and cells with a tweezer, and after centrifugation at 1,200 rpm for 5 minutes the supernatant is discarded. Then the pellet is
5 treated with Tris-HCl buffer (pH 7.65) for 1 to 2 minutes, and erythrocytes are eliminated. After washing 3 times with MEM, spleen cells used for cell fusion are obtained.

[0026]

(2) Preparation of permanently proliferating cells

10 Any cell with permanent proliferation ability can be used as the permanently proliferating cell for the fusion. However, myeloma cells are generally used. It is preferable to use cells derived from the same specie as that from which the antibody-producing cell was obtained. For example, in the case of mouse, the following cell
15 lines are known as 8-azaguanine resistant mouse (BALB/c)-derived myeloma cell lines.

P3-X63Ag8-U1 (P3-U1) (Current.Topics in Microbiol.Immunol., 81:1-7, (1978)),

P3/NS1/1-Ag4-1 (NS-1) (Eur.J.Immunol., 6:511-519, (1976)),

20 SP2/0-Ag14 (SP-2) (Nature, 276:269-270, (1978)),

P3-X63-Ag8653 (653) (J.Immunol., 123:1548-1550, (1979), and

P3-X63-Ag8 (X63) (Nature, 256:495-497, (1975))

These permanently proliferating cell lines are subcultured in 8-azaguanine media (normal media, RPMI-1640 medium supplemented with
25 glutamine (1.5 mM), 2-mercaptoethanol (5×10^{-5} M), gentamicin (10 μ g/mL) and fetal calf serum (FCS, CLS) (10%), supplemented with 8-azaguanine (15 μ g/mL)), and passaged to normal media 3 to 4 days before cell fusion to ensure a cell count of 2×10^7 cells at the day of cell fusion.

30 [0027]

(3) Cell fusion

Fusion of cells is, for example, performed as follows. The antibody-producing cell obtained in (1), and the permanently
35 proliferating cell prepared in (2) are mixed at a cell rate of 5 to 10:1 after washing well with MEM or PBS. After centrifugation at 1,200 rpm for 5 minutes, the supernatant is removed and the precipitated

cells are detached well. 0.1 to 1.0 mL/ 10^8 cells are added to a mixture of 1 to 4 g polyethylene glycol-1000 (PEG-1000), 1 to 4 mL MEM, and 0.5 to 1.0 mL dimethyl sulfoxide, and the mixture is kept at 37°C while stirring to fuse the cells. Then, the mixture is diluted by adding 3 mL MEM few times every 10 minutes to a total volume of 50 mL MEM to quench the cell fusion reaction. Next, the supernatant is discarded after centrifugation (1,500 X 5 min), cells are gently tapped, 100 mL normal medium (RPMI-1640 media, 10% FCS) is added, and the cells are suspended by gentle pipetting with a measuring pipette. 100 µL suspension is dispensed to each well of a 96 well culture plate, and incubated at 37°C for 3 to 5 days in an 5% CO₂ incubator. 100 µL of HAT medium (normal medium supplemented with hypoxanthine (10^{-4} M), thymidine (1.5×10^{-5} M), and aminopterin (4×10^{-7} M)) is added to each well of the culture plate, and incubated further for 3 days. Thereafter, half of the culture supernatant is discarded and the same amount of HAT medium is added thereto every three days. The incubation is continued for about 2 weeks at 37°C in a 5% CO₂ incubator.

[0028]

(4) Screening and cloning of the hybridoma

Supernatants are removed from wells in which fusion cells are observed to grow in colonies, same amount of HT medium (HAT medium without aminopterin) is added, and the cells are incubated for 4 days. Part of the culture supernatant is harvested to measure the antibody titer against megsin protein by the aforementioned protein A rosette assay. The antibody titer can be also assayed by other methods like the following. For example, 1) by adding the hybridoma culture supernatant to a solid phase (e.g. microplate), where the megsin protein antigen is attached directly or together with a carrier; 2) adding anti-immunoglobulin antibody (if the cell for cell fusion is derived from mouse, anti-mouse immunoglobulin antibody is used) labeled with radioactive substances or enzymes, or protein A; and 3) detecting the anti-megsin protein monoclonal antibody bound to the solid phase. The antibody titer can be also assayed by: 1) adding the hybridoma culture supernatant to a solid phase, where an anti-immunoglobulin antibody or protein A is attached; 2) adding

megsin protein labeled with radioactive substances or enzymes, and
3) detecting the anti-megsin protein monoclonal antibody bound to
the solid phase.

Cloning is repeated 4 times by limiting dilution method for
5 wells in which the production of antibodies that react to the mesgin
protein was observed, and cell lines, which show stable mesgin protein
antibody titers are selected as anti-megsin protein monoclonal
antibody producing hybridoma cell lines.

[0029]

10 (5) Preparation of the monoclonal antibody

The monoclonal antibody is produced by incubating the thus
obtained hybridomas *in vitro* or *in vivo*. In case where the incubation
is conducted *in vivo*, the hybridoma is transplanted to appropriate
animals. It is preferable to use the same kind of animal as that from
15 which the spleen cells for cell fusion were taken. For example, 2
to 4 X 10⁶ cells of anti-megsin protein monoclonal antibody producing
hybridoma cells obtained in (4) is administered intraperitoneally
to each 8 to 10 week old BALB/c female mouse treated with pristane
(reared for 2 weeks after intraperitoneal administration of 0.5 mL
20 2,6,10,14-tetramethylpentadecane-pristane). Subsequently, after 2
to 3 weeks, the abdomen of the mouse enlarges due to the accumulation
of ascites, which contain the monoclonal antibody at a high
concentration, in the visceral cavity of the mouse. The ascites are
then collected from the mouse, centrifuged (3,000 rpm X 5 min) to
25 remove solid matters, and the IgGs are purified. Alternatively, the
hybridoma is cultured *in vitro* in a serum-free and an appropriate
amount of antibody secretes to the supernatant. The ascites or the
culture supernatant is salted out with 50% ammonium sulfate, and then
dialyzed for 1 to 2 weeks with PBS. The dialyzed fraction is passed
30 through a protein A Sepharose column, and the IgG fraction is collected
to obtain a purified monoclonal antibody.

The isotype of the antibody is determined according to the
Ouchterlony method (double immunodiffusion) ("Menekigaku jikken
nyumon, Seibutsugaku jikkenhou 15" (A laboratory manual for immunology,
35 Biochemical experimentation 15), Gakkai Shuppan Center, p74 (1981)).
The quantity of the protein is calculated by the Folin method and

by the absorbance at 280 nm ($1.4 \text{ (OD}_{280}) \div \text{immunoglobulin } 1 \text{ mg/mL}$).

[0030]

(6) Characterization of the monoclonal antibody

The characterization of the monoclonal antibody obtained above
5 can be performed by, for example, (1) immunoprecipitation method
utilizing human lymphocyte derived cell lines, such as HSB-2, K562,
and so on, the cell surface of which is labeled with iodine (J.Immunol.,
138: 2850-3855 (1987)); (2) enzyme immunoassay (ELIZA) (J.Immunol.,
142: 2743-2750 (1989)); and such.

10 [0031]

(7) Preparation of labeled monoclonal antibody

The obtained purified monoclonal antibody may be enzyme labeled
by methods like glutaraldehyde method (Immunochem., 6: 43 (1969)),
periodic acid method (J.Histochem.Cytochem., 22: 1084 (1974)),
15 maleimide method (J.Biochem., 79: 233 (1976)), pyridil disulfide
method (Biochem.J., 173: 723 (1978)), etc.

For example, the periodic acid method is performed as follows:

(1) periodic acid (38.5 mg/mL) is added to the peroxidase solution
(4 mg/mL) while stirring; (2) reacted at room temperature for 20
20 minutes; (3) the buffer is exchanged using PD-10 (Amersham Pharmacia
Biotech) substituted with 1 mM acetate buffer (pH 4.5); (4) 40 μL
of 0.2 M sodium hydroxide is added; (5) 10 mg of monoclonal antibody
dialyzed with 10 mM carbonate buffer is added (pH 9.5); (6) react at
room temperature for 2 hours; (7) after the reaction is completed,
25 the mixture is cooled on ice; (8) 100 μL solution of sodium boron
hydride (4 mg/mL) is added and reacted for 2 hours; (9) the reaction
solution is changed to PBS using PD-10; (10) centrifuged at 3,000
rpm for 30 minutes; (11) the supernatant is gel filtrated using
Sephacryl S200HR26 X 30 (Amersham Pharmacia Biotech); (12) the labeled
30 monoclonal antibody fraction is obtained by detecting the absorbance
at 403 and 280 nm; (13) bovine serum albumin (10 mg/mL) is added to
the obtained fraction; (14) stored at -20°C , and diluted with
PBS-Tween (Trademark) 20 before usage.

[0032]

35 (B) Preparation of granules for detection

The granules for detection used herein may be prepared by

physically or chemically binding anti-megsin protein monoclonal antibody to appropriate granules, for example, gel for chromatography. The method used in the present invention for binding the antibody to chemically activated granules is a preferable method due to the expected binding stability. More specifically, the method for binding the antibody used in the present invention to granules that are actively tosylated by p-toluenesulfonyl chloride can be given.

Granules to be used are made of glass, agarose, Sepharose, agarose filled porous diatomite, hydrophilic copolymerized acrylic gel, polystyrene, and so on. Use of magnetic granules enables collection of the granules with the use of magnet, and so on, thereby allowing detection of small quantity samples. For example, by placing substances capable of being magnetic (e.g. Fe_2O_3) in the core of the granules, it is possible to make the granules superparamagnetic. Such granules are commercially available as solid phases for immunological analyses. The granule may take any appropriate form such as a sphere, amorphous fracture, etc., however, sphere shaped granules are preferred. There is no limitation on the particle size of the granule, and an average particle size, for example, of 5 to 1000 μm can be given. Furthermore, the use of granules with larger specific gravity than that (about 1) of the reaction solution facilitates collection of the granules, giving an effect similar to that when magnetic granules are used. Furthermore, the use of granules with larger specific gravity is more advantageous for antibodies whose binding is easily disassociated, because gentle centrifugation conditions are possible when collecting the granules.

Anti-megsin protein antibodies can be bound to the granules not only by direct binding but also by indirect binding. For example, if a mouse monoclonal antibody is used as the antibody, an antibody that recognizes mouse IgG is attached to the granule and the mouse antibody can be bound to the granule indirectly. Such antibodies are called secondary antibodies. Indirect binding can be also accomplished by utilizing protein A or protein G, which bind to the constant region of immunoglobulin, or by capturing avidin-attached granules with biotinylated antibodies, and so on. To chemically bind the secondary antibody, protein A, or protein G to the granule, it

is preferable to activate the granules before binding. As an activation method of the granule, any appropriate activation method used to conjugate proteins to the granule can be selected. Such activation methods include, the tosyl chloride method, bromocyanide method, bromoacetyl method, glutaraldehyde method, etc. There are also commercially available activated granules. Activation and binding of these activated granules to proteins, such as secondary antibodies, protein A, protein G, and such, can be performed according to conventional methods.

[0033]

In addition, granules conjugated to secondary antibodies, protein A, protein G, and so on are also commercially available. Granules mentioned below are known as commercially available granules.

Dynabeads™ M-450, M-280 provided by Veritas, and imported by Dynal Japan

Sheep anti-mouse IgG coated type

Goat anti-mouse IgG coated type

Sheep anti-rat IgG coated type

Sheep anti-rabbit IgG coated type

Polyscience Inc.

Goat anti-mouse IgG (H&L) carboxylate beads

Goat anti-rabbit IgG (H&L) carboxylate beads

Protein A carboxylate beads

Goat anti-rabbit IgG (H&L) micromagnet particle

Goat anti-rabbit IgG (H&L) micromagnet particle

Protein A micromagnet particle

Sheep anti-mouse IgG (H&L) micromagnet particle

The binding of the present antibody to the granules above is conducted as follows: (1) unspecific binding is prevented by treating the granules suspended in appropriate media with a protein solution, and (2) mixing the ascites containing the antibody or the purified antibody solution.

[0034]

(C) Detection method

The detection method of the present invention is accomplished

by collecting blood or urine from the subject, and using the supernatant after centrifugation as the specimen. The centrifugation of urine is done to separate the precipitate, and the supernatant of urine left standing still, or urine from which the precipitate is removed by filtration can be also used. Specimen diluted with the detection granule prepared above and the labeled antibody obtained in (7) are mixed, and then, incubated at room temperature for 2 hours. After the reaction has come to an end, the mixture is washed, and the substrate solution is added for chromogenic reaction. Then, the mixture is centrifuged to remove the granules, the supernatant is transferred onto a microplate, and the absorbance is measured. A specimen from a normal healthy person is also assayed according to the same method, and the absorbance is compared. Mere comparison of the concentration of the megsin protein, as well as a comparison of the absolute quantity of megsin protein in the body fluid, which is obtained by multiplying with the volume of body fluid of the individual, or comparisons based on of other similar corrected values might be performed.

[0035]

20 (D) Kit

The materials used to perform the experiment above can be supplied as a kit. The kit may comprise a detection granule, to which the antibody is conjugated as described above, and a magnet. It may also include an antibody labeled with a marker molecule. Further, the kit of the present invention may include test tubes, centrifugation tubes, other similar containers, pipettes or other suction instruments, and microscopes. Furthermore, enzyme substrates necessary for detecting the label, and positive or negative standard samples may be combined. Solid granules and antibodies, which are the raw material for the detection granules mentioned above, may be included instead of the detection granules. The invention is described in detail with reference to the examples below, but it is not to be construed as being limited thereto.

[0036]

35 [Examples]

[Example 1] Production of polyclonal antibodies against synthetic

peptides of the megsin protein

Polyclonal antibodies against megsin protein were produced by using regions having low identity with other members of the serpin family and which are hydrophilic as the immunogens. Peptides having
 5 the amino acid sequences shown below with a cystein attached to the N-terminus or the C-terminus were synthesized by solid phase methods.

Position from	Amino acid sequence	SEQ ID NO/No.
N-terminus		
16- 30	CFREMDDNQGNVFF	11/1
72- 86	CSQSGLQSQLKRVFSD	12/2
339-354	ATGSNIVEKQLPQSTLC	13/3
342-356	CSNIVEKQLPQSTLFR	14/342

Each of these synthetic peptides were bound to bovine
 10 thyroglobulin (Sigma) using N-(6-maleimidocaproyloxy)succinimide (Dojindo Molecular Technologies Inc.), were dialyzed against 0.85% NaCl solution, and were mixed well with the adjuvant (Difco) to emulsion, and were administered subcutaneously to a rabbit. 3 weeks after priming (20 µg/rabbit), second immunization (50 µg/rabbit) was
 15 carried out, and thereafter, four immunizations (50, 100, 200 µg/rabbit) were performed every two weeks. Freund's complete adjuvant was used for the priming, and thereafter, incomplete Freund's adjuvant was used. After 41 days and 55 days, the antibody titer of the serum was evaluated by enzyme-linked immunosorbent assay (ELISA)
 20 to confirm that the serum obtained by blood sampling reacts with synthetic peptide. To conduct primary reaction, 100 µL of continuously diluted antiserum was added to each well of the 96 well plate, in which 50 ng/well antigen had been immobilized. After washing, the HRP conjugated goat anti-rabbit IgG (Cappel) was reacted
 25 against it in the second reaction. After washing, the coloring reaction was done using orthophenylenediamine (Wako Pure Chemical Industries, Ltd.) as the substrate and absorbance was measured at 492 nm. An increase in antibody titer was confirmed.

[0037]

30 [Example 2] Purification of polyclonal antibodies against synthetic

peptides of megsin protein

The polyclonal antibodies against each synthetic peptide of megsin protein were purified by immunoaffinity chromatography according to conventional methods (Cell Engineering supplement 5 "Jikken Protocol Series Kou-peptide Jikken protocol (Experimental protocol series: Anti-peptide experimental protocol)", Shujun-sha, Japan). More specifically, the method was carried out as follows: 1) an affinity column was prepared by immobilizing each synthetic peptide to FMP (2-fluoro-1-methylpyridinium toluene-4-sulfonate) 10 activated Cellulofine (Seikagaku Corp.); 2) after the rabbit serum showing increased antibody titer was diluted with PBS (-), the antibody was purified by affinity using the peptide column. It was demonstrated that the obtained purified antibody is specific to the megsin protein, by confirming its reaction against megsin protein 15 fusion protein by Western blotting (Fig. 1 to 3).

[0038]

[Example 3] Studies on reactivity of the rabbit polyclonal anti-megsin peptide IgG

The reactivity of the rabbit IgG, which was produced using 20 megsin protein as the immunogen, was studied using the following various proteins as antigens. MBP-megsin protein and His-megsin protein were prepared as follows: 1) the coding region was amplified by PCR based on the nucleotide sequence of megsin gene described in SEQ ID NO: 1; 2) the amplified products were inserted into maltose 25 binding protein fusion protein expression vector pMAL-c (New England Biolab) to obtain vectors that express the fusion protein of MBP and megsin protein; 3) E.coli were transformed with this vector; and then, 4) the MBP-megsin protein was obtained by purifying the expression product from the disrupted cell solution by affinity chromatography 30 utilizing amylose, and such. To obtain His-megsin protein, a fusion protein expression vector containing aforementioned megsin gene in the vector ptrcHisA (Invitrogen), which expresses the inserted gene as a fusion protein with a histidine tag, was constructed. This vector was transformed into E.coli JM109, and following the induction of 35 expression by IPTG, the His-megsin protein was obtained as a fusion protein purified by nickel column and such from the disrupted cell

solution.

The megsin protein(+)virus-infected silkworm body fluids refers to body fluids of a silkworm, which is infected with megsin gene introduced virus according to known methods (J.General Virology 5 78: 3073-3080 (1979)). On the other hand, megsin protein(-)virus-infected silkworm body fluids refers to body fluids of a silkworm, which is infected with a virus without the megsin gene. The gene encoding megsin necessary for designing these megsin recombinants can be synthesized by PCR using mRNA of the mesangial 10 cell as the template, or obtained from the megsin gene cloning vector (Accession No. FERM BP-6518) constructed by the inventors.

Reactivity against plasminogen activator inhibitor-1 (PAI I), MBP-plasminogen activator inhibitor-2 fusion protein (MBP-PAI II), and so on was also observed. These proteins were selected to confirm 15 the cross-reactivity of the antibody, and share some extent of homology with the megsin protein.

MBP-megsin protein,
His-megsin protein,
MBP-PAI II,
20 PAI I,
megsin protein expressed in CHO cells,
megsin protein (+) virus-infected silkworm body fluid,
megsin protein (-)virus-infected silkworm body fluid,
non-virus-infected silkworm body fluid, and
25 human serum albumin.

[0039]

Each protein solution was treated with same amount of sample buffer (0.25% Tris-HCl, 2% SDS, 30% glycerin, 10% β -mercaptoethanol, 0.025% Bromophenol Blue) (Daiichi Pure Chemicals), heated at 100°C 30 for 5 minutes, and the proteins were separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (Laemmli, U.K., Nature 227: 680-685 (1970)). The proteins separated by SDS-PAGE was blotted on poly vinylidene difluoride (PVDF) membrane (BioRad) with a constant voltage of 100 V for 1 hour, using the blotting solution (25mM Tris-HCl, 35 192 mM glycine, 20% methanol, pH 8.3). Following the washing of the blotted PVDF membrane with distilled water, the membrane was blocked

for 3 hours in TTBS solution containing 5% Block Ace. After washing the PVDF membrane with TTBS (20 mM Tris, 500 mM NaCl, 0.05% Tween20, pH 7.5), it was reacted at 4°C, overnight with the primary antibody, rabbit polyclonal anti-megsin peptide IgG, diluted in TTBS. Then it was detected with Amplified Alkaline Phosphatase Immuneblot Kit (BioRad). That is, it was first incubated for 1 hour with TTBS diluted biotinylated goat anti-rabbit IgG at room temperature, then reacted with streptavidin - biotinylated alkaline phosphatase complex, which was prepared in advance by incubating streptavidin and biotinylated alkaline phosphatase for 1 hour at room temperature. The PVDF membrane was washed in TTBS, and then incubated with the substrate (nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl phosphate, p-toluidine salt solution) for about 30 minutes at room temperature to visualize the antibody bound to the primary antibody. The reaction was quenched by reacting with a sufficient amount of distilled water.

[0040]

[Example 4] Enzyme labeling of affinity purified anti-megsin peptide IgG

The affinity purified anti-megsin peptide IgG obtained in Example 2 was labeled with alkaline phosphatase using a commercial labeling kit (AP labeling kit: Boehringer-Mannheim) following the instructions in the kit. Biotinylation was performed with another labeling kit (ECL protein biotinylation module: Amersham Pharmacia Biotech) following the instructions.

[0041]

[Example 5] Measurement of urinary megsin protein by ELISA

Urine from 16 normal healthy persons, 24 IgA nephropathy patients, and 24 minimal-change nephritic syndrome patients were centrifuged to exclude the insoluble material. This was then concentrated using an ultrafilter, with a molecular weight cut-off of 5,000 (Ultrarafree, Millipore). Each well of the 96 well ELISA plate was coated with rabbit polyclonal anti-megsin peptide-2 IgG. 120 µL PBS (-) was poured into each well, then 120 µL concentrated urine was added thereto and was left standing overnight at 4°C. The wells were washed with PBS (-), and Block Ace (Dainippon Pharmaceutical) was added to conduct blocking for 2 hours at room temperature. The wells

were then washed with PBS (-) containing 0.05%(w/v) Tween20 (Wako Pure Chemical), alkaline phosphatase labeled rabbit polyclonal anti-megsin peptide-1 antibodies were added, and left standing for an hour at room temperature. After washing with Tween-PBS, 100 μ L of coloring substrate solution (1 mg/mL orthophenylenediamine (Wako Pure Chemical) - 1 M diethanolamine (Wako Pure Chemical) buffer (pH 9.8)) was added to each well. After reacting at room temperature, the reaction was quenched with 50 μ L/well 1 N sodium hydroxide, and the absorbance at 405nm was measured with a microplate reader (Spectra MAX 250, Molecular Devices). The megsin protein concentration in the urine was determined from the calibration curve. The creatinine concentration in the same urine specimen was determined with the commercially available creatinine-measuring reagent (measuring reagent for automated analyzer "Dia"-Crea, Dia Pharmaceutical), and the amount of megsin protein was corrected as creatinine ratio. The result is shown in Fig. 4. A significant difference in megsin concentration was observed between that of the normal healthy persons and patients of IgA nephropathy ($p < 0.001$), or between that of IgA nephropathy patients and minimal-change nephritic syndrome patients ($p < 0.001$).

Since a significant difference between normal healthy persons and IgA nephropathy patients was observed, onset of renal disorders could be confirmed by measuring urinal megsin protein. Furthermore, due to the significant difference observed between patients of IgA nephropathy and minimal-change nephritic syndrome, it was revealed that these diseases could be distinguished by measuring urinal megsin protein according to the present invention. These diseases cannot be distinguished by conventional screening methods using urinary protein as indicators, because positive results would be shown for both diseases.

Furthermore, in comparisons of urinary megsin protein concentration without creatinine correction, although megsin protein concentration tends to be higher for patients with renal disorders, it seemed difficult to prove a statistically significant difference with a small number of examples. Thus, it is preferable to compare the amounts by methods like creatinine correction and such, when

implementing the measurement of megsin protein in urine according to the present invention.

[0042]

[Example 6] Detection and assay of urinary megsin protein by Western blotting

Urine from 8 normal healthy persons and 9 IgA nephropathy patients were centrifuged, after discarding insoluble material, the solution was concentrated with an ultrafilter with a molecular weight cut-off of 5,000 (Ultrafree, Millipore). The concentrated urine was treated with same amount of sample buffer (0.25% Tris-HCl, 2% SDS, 30% glycerin, 10% β -mercaptoethanol, 0.025% Bromophenol Blue) (Daiichi Pure Chemicals), heated at 100°C for 5 minutes, and separated in a gradient gel with a gel gradient of 4 to 20% (Daiichi Pure Chemicals) by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (Laemmli, U.K. (1970) Nature 227, 680-685). The protein separated by SDS-PAGE was blotted on poly vinylidene difluoride (PVDF) membrane (BioRad) with constant voltage of 100 V for 1 hour, using the blotting solution (25 mM Tris-HCl, 192 mM glycine, 20% methanol, pH 8.3). Following the washing of the blotted PVDF membrane with distilled water, the membrane was blocked for 3 hours in TTBS solution containing 5% Block Ace. After washing the PVDF membrane with TTBS (20 mM Tris, 500 mM NaCl, 0.05% Tween20, pH 7.5), it was reacted overnight at 4°C with a primary antibody solution, rabbit polyclonal anti-megsin peptide IgG diluted in TTBS.

[0043]

Then it was detected with Amplified Alkaline Phosphatase Immuneblot Kit (BioRad). That is, it was first incubated for 1 hour with biotinylated goat anti-rabbit IgG diluted with TTBS at room temperature, then reacted with streptavidin-biotinylated alkaline phosphatase complex, which was prepared in advance by incubating streptavidin and biotinylated alkaline phosphatase for 1 hour at room temperature. The PVDF membrane was washed in TTBS, and then incubated with the substrate (nitro blue tetrazolium and 5-bromo-4-chloro-3-indolyl phosphate, p-toluidine salt solution) for about 30 minutes at room temperature to visualize the antibody bound to the primary antibody. The reaction was quenched by reacting

with a sufficient amount of distilled water. The visualized bands were quantitatively analyzed using Densitograph gel documentation system (AE-6920M-05, Atto) and densitograph analyzing software (AE-6920WSE/MSE, Atto). The result (electrophoretogram) is shown in Fig. 5. The measurements of the image analyzer are as follows. Dense bands were observed for IgA nephropathy patients compared to the bands of concentrated urine of normal healthy persons. It was revealed that quantitative measurement of the megsin protein is possible by quantitatively analyzing the density of these bands.

10 Measurement result of the image analyzer

Lane	Measurement result
1	Molecular weight marker
2	8.5
3	5.2
4	13.8
5	4.8
6	32.8
7	15.1

[0044]

[Example 7] Preparation of granules for detection

The granules for detection of the present invention were produced by conjugating anti-megsin peptide IgG to commercially available magnetic granules through anti-mouse IgG antibody. Dynabeads™ M-450 tosylactivated (homogenous tosyl activated paramagnetic polystyrene granules: Dynal Biotech) was used as the granules of the present invention, and a 1mM hydrochloric acid solution thereof was washed once with sterile distilled water and then with 0.2 M boric acid buffer (pH9.5). Then, anti-megsin peptide IgG antibody was dissolved in 0.2 M boric acid buffer (pH 9.5) at a concentration of 150 µg/mL, and this solution was added to an equal amount of the activated granule suspension (antibody/granule=75 µg/15mg). Then, the mixture was incubated for 24 hours at 22°C with gentle stirring. The antibody-conjugated granules were collected with a magnet, and the supernatant was discarded leaving the granules attached to the magnet. Then the granules were washed as follows:

- (1) with 5 mL 0.1 M PBS (pH7.0) for 10 minutes,
- (2) with 5 mL 1 M ethanolamine-HCl (pH 9.5) for 2 hours,
- (3) with 5 mL 0.05 M Tris (pH 7.5) containing 0.1 M sodium chloride, 0.1% bovine serum albumin (BSA), and 0.1% Tween20 for 12 hours, and
- 5 (4) with 5 mL buffer of (3) without Tween20 for 2 hours.

Thereafter, granules were collected with a magnet, supernatant was discarded, and the granules were suspended in PBS (-)/BSA containing 10% Block Ace at a concentration of about 4×10^8 granules/mL (30 mg/mL). Thus obtained IgG-conjugated granules are stable for at least 6 months
10 at 4°C.

[0045]

[Example 8] Measurement of urinary megsin protein by the magnetic granule method

Megsin protein concentration of occasional urine samples from
15 12 normal healthy persons and 24 patients with different renal diseases were measured by the magnetic granule method utilizing antibody-conjugated granules prepared in Example 7. 12 of the renal disease patients were affected with IgA nephropathy, and the other 12 were affected with minimal-change nephritic syndrome. The urinary
20 samples were kept at -80°C, and were thawed before measurement. They were centrifuged at 3,000 rpm for 5 minutes, and the supernatant thereof were used as the sample. The measurement was carried out as follows:

500,000 magnetic granules conjugated to the primary antibody
25 (rabbit polyclonal anti-megsin peptide-2 antibody) were put into a tube (1.5 mL) blocked beforehand with Block Ace (Dainippon Pharmaceutical). 500 µL of each urinary sample and equal amounts of alkaline phosphatase labeled rabbit polyclonal anti-megsin peptide-1 antibodies dialyzed with PBS (-) were mixed, and reacted for 2 hours
30 at room temperature while rotating. Then, after centrifugation at 3,000 rpm for 5 minutes, leaving the granules attached to the magnet, the supernatant was discarded, and 1mL washing solution (0.05% (w/v) Tween 20 (Wako Pure Chemical) containing PBS (-) (pH7.4) (Tween-PBS)) was added and the granules were washed while stirring. The washing
35 was repeated 4 times, and then, 100 µL substrate solution was added to react for 20 to 30 minutes with stirring. 1 M diethanolamine (Wako

Pure Chemical) buffer (pH 9.8) containing 1 mg/mL orthophenylenediamine (Wako Pure Chemical) was used as the substrate solution. Following the completion of the reaction, the reaction was quenched with 50 μ L 1 N sodium hydroxide, and absorbance at 405 nm was measured. The urinary megsin protein concentration was calculated from the calibration curve. Creatinine concentration of the same urinary sample was measured as in Example 5, and the megsin protein concentration was corrected as creatinine ratio.

According to the result of the measurement, urinary megsin protein measurement values (creatinine correction value: AU) of IgA nephropathy patients were distributed in the top region compared to those of the normal healthy person. Additionally, the urinary megsin protein measurement values of IgA nephropathy patients were also higher than those of minimal-change nephritic syndrome patients. Thus, according to the present invention, it is possible not only to decide whether a person is affected by a renal disorder, but definitely identify the state of the renal disorder.

[0046]

[Effects of the Invention]

The present invention facilitates the evaluation of renal functions not by a highly invasive test like renal biopsy, but by using invasive samples like urine and blood. The realization of a simple *in vitro* evaluation of renal functions enables a quick and inexpensive test service that can be provided to many more subjects than before. There are many renal diseases, in which it is better to delay the initiation of dialysis treatment by an early detection of the diseases. Thus, the present invention not only eases the burden on the subjects, but also enables detection and observation of more patients who would need dialysis treatment in the future by providing methods for evaluating renal functions with simple examination procedures. Furthermore, a high evaluation reliability can be expected thanks to the excellent indicator, the megsin protein found in biological specimens like urine and blood, which reflects renal function.

Particularly, the immunological detection method of the megsin protein, and the kit used for the method, which are provided by the

present invention, are superior in manipulatability utilizing magnetic granule carriers. Quick and accurate measurement of the megsin protein is enabled by the present detection method. Furthermore, since the method enables the understanding of the degree of mesangial cell proliferation, the present invention makes it easy to determine the timing of when to commence or whether or not to continue steroid therapy used for treating IgA nephropathy.

[0047]

[Sequence Listing]

SEQUENCE LISTING

<110> MIYATA, Toshio

KUROKAWA, Kiyoshi

Fuso Pharmaceutical Industries, Ltd.

<120> A method of Detecting the Megsin Protein, and Use Thereof.

<130> F2-101

<140>

<141>

<160> 18

<170> PatentIn Ver. 2.0

<210> 1

<211> 1143

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1).. (1140)

<300>

<302> A mesangium-predominant gene, megsin, is a new serpin
upregulated in IgA nephropathy.

<303> J. Clin. Invest.

5 <304> 120

<305> 4

<306> 828-836

<307> 1998-8-15

10 <400> 1

atg gcc tcc ctt gct gca gca aat gca gag ttt tgc ttc aac ctg ttc 48
Met Ala Ser Leu Ala Ala Ala Asn Ala Glu Phe Cys Phe Asn Leu Phe
1 5 10 15

15 aga gag atg gat gac aat caa gga aat gga aat gtg ttc ttt tcc tct 96
Arg Glu Met Asp Asp Asn Gln Gly Asn Gly Asn Val Phe Phe Ser Ser
20 25 30

ctg agc ctc ttc gct gcc ctg gcc ctg gtc cgc ttg ggc gct caa gat 144
20 Leu Ser Leu Phe Ala Ala Leu Ala Leu Val Arg Leu Gly Ala Gln Asp
35 40 45

gac tcc ctc tct cag att gat aag ttg ctt cat gtt aac act gcc tca 192
25 Asp Ser Leu Ser Gln Ile Asp Lys Leu Leu His Val Asn Thr Ala Ser
50 55 60

gga tat gga aac tct tct aat agt cag tca ggg ctc cag tct caa ctg 240
Gly Tyr Gly Asn Ser Ser Asn Ser Gln Ser Gly Leu Gln Ser Gln Leu
65 70 75 80

30 aaa aga gtt ttt tct gat ata aat gca tcc cac aag gat tat gat ctc 288
Lys Arg Val Phe Ser Asp Ile Asn Ala Ser His Lys Asp Tyr Asp Leu
85 90 95

35 agc att gtg aat ggg ctt ttt gct gaa aaa gtg tat ggc ttt cat aag 336
Ser Ile Val Asn Gly Leu Phe Ala Glu Lys Val Tyr Gly Phe His Lys

	100	105	110	
	gac tac att gag tgt gcc gaa aaa tta tac gat gcc aaa gtg gag cga	384		
	Asp Tyr Ile Glu Cys Ala Glu Lys Leu Tyr Asp Ala Lys Val Glu Arg			
5	115	120	125	
	ggt gac ttt acg aat cat tta gaa gac act aga cgt aat att aat aag	432		
	Val Asp Phe Thr Asn His Leu Glu Asp Thr Arg Arg Asn Ile Asn Lys			
	130	135	140	
10	tgg gtt gaa aat gaa aca cat ggc aaa atc aag aac gtg att ggt gaa	480		
	Trp Val Glu Asn Glu Thr His Gly Lys Ile Lys Asn Val Ile Gly Glu			
	145	150	155	160
15	ggt ggc ata agc tca tct gct gta atg gtg ctg gtg aat gct gtg tac	528		
	Gly Gly Ile Ser Ser Ser Ala Val Met Val Leu Val Asn Ala Val Tyr			
	165	170	175	
	ttc aaa ggc aag tgg caa tca gcc ttc acc aag agc gaa acc ata aat	576		
20	Phe Lys Gly Lys Trp Gln Ser Ala Phe Thr Lys Ser Glu Thr Ile Asn			
	180	185	190	
	tgc cat ttc aaa tct ccc aag tgc tct ggg aag gca gtc gcc atg atg	624		
	Cys His Phe Lys Ser Pro Lys Cys Ser Gly Lys Ala Val Ala Met Met			
25	195	200	205	
	cat cag gaa cgg aag ttc aat ttg tct gtt att gag gac cca tca atg	672		
	His Gln Glu Arg Lys Phe Asn Leu Ser Val Ile Glu Asp Pro Ser Met			
	210	215	220	
30	aag att ctt gag ctc aga tac aat ggt ggc ata aac atg tac gtt ctg	720		
	Lys Ile Leu Glu Leu Arg Tyr Asn Gly Gly Ile Asn Met Tyr Val Leu			
	225	230	235	240
35	ctg cct gag aat gac ctc tct gaa att gaa aac aaa ctg acc ttt cag	768		
	Leu Pro Glu Asn Asp Leu Ser Glu Ile Glu Asn Lys Leu Thr Phe Gln			

		245		250		255	
		aat cta atg gaa tgg acc aat cca agg cga atg acc tct aag tat gtt	816				
		Asn Leu Met Glu Trp Thr Asn Pro Arg Arg Met Thr Ser Lys Tyr Val					
5		260 265 270					
		gag gta ttt ttt cct cag ttc aag ata gag aag aat tat gaa atg aaa	864				
		Glu Val Phe Phe Pro Gln Phe Lys Ile Glu Lys Asn Tyr Glu Met Lys					
		275 280 285					
10		caa tat ttg aga gcc cta ggg ctg aaa gat atc ttt gat gaa tcc aaa	912				
		Gln Tyr Leu Arg Ala Leu Gly Leu Lys Asp Ile Phe Asp Glu Ser Lys					
		290 295 300					
15		gca gat ctc tct ggg att gct tcg ggg ggt cgt ctg tat ata tca agg	960				
		Ala Asp Leu Ser Gly Ile Ala Ser Gly Gly Arg Leu Tyr Ile Ser Arg					
		305 310 315 320					
		atg atg cac aaa tct tac ata gag gtc act gag gag ggc acc gag gct	1008				
20		Met Met His Lys Ser Tyr Ile Glu Val Thr Glu Glu Gly Thr Glu Ala					
		325 330 335					
		act gct gcc aca gga agt aat att gta gaa aag caa ctc cct cag tcc	1056				
		Thr Ala Ala Thr Gly Ser Asn Ile Val Glu Lys Gln Leu Pro Gln Ser					
25		340 345 350					
		acg ctg ttt aga gct gac cac cca ttc cta ttt gtt atc agg aag gat	1104				
		Thr Leu Phe Arg Ala Asp His Pro Phe Leu Phe Val Ile Arg Lys Asp					
		355 360 365					
30		gac atc atc tta ttc agt ggc aaa gtt tct tgc cct tga	1143				
		Asp Ile Ile Leu Phe Ser Gly Lys Val Ser Cys Pro					
		370 375 380					
35							

<211> 380

<212> PRT

<213> Homo sapiens

5 <400> 2

Met Ala Ser Leu Ala Ala Ala Asn Ala Glu Phe Cys Phe Asn Leu Phe
 1 5 10 15

Arg Glu Met Asp Asp Asn Gln Gly Asn Gly Asn Val Phe Phe Ser Ser
 10 20 25 30

Leu Ser Leu Phe Ala Ala Leu Ala Leu Val Arg Leu Gly Ala Gln Asp
 35 40 45

Asp Ser Leu Ser Gln Ile Asp Lys Leu Leu His Val Asn Thr Ala Ser
 50 55 60

Gly Tyr Gly Asn Ser Ser Asn Ser Gln Ser Gly Leu Gln Ser Gln Leu
 65 70 75 80

Lys Arg Val Phe Ser Asp Ile Asn Ala Ser His Lys Asp Tyr Asp Leu
 85 90 95

Ser Ile Val Asn Gly Leu Phe Ala Glu Lys Val Tyr Gly Phe His Lys
 25 100 105 110

Asp Tyr Ile Glu Cys Ala Glu Lys Leu Tyr Asp Ala Lys Val Glu Arg
 115 120 125

Val Asp Phe Thr Asn His Leu Glu Asp Thr Arg Arg Asn Ile Asn Lys
 130 135 140

Trp Val Glu Asn Glu Thr His Gly Lys Ile Lys Asn Val Ile Gly Glu
 145 150 155 160

Gly Gly Ile Ser Ser Ser Ala Val Met Val Leu Val Asn Ala Val Tyr

	165	170	175
	Phe Lys Gly Lys Trp Gln Ser Ala Phe Thr Lys Ser Glu Thr Ile Asn		
	180	185	190
5	Cys His Phe Lys Ser Pro Lys Cys Ser Gly Lys Ala Val Ala Met Met		
	195	200	205
	His Gln Glu Arg Lys Phe Asn Leu Ser Val Ile Glu Asp Pro Ser Met		
10	210	215	220
	Lys Ile Leu Glu Leu Arg Tyr Asn Gly Gly Ile Asn Met Tyr Val Leu		
	225	230	235 240
15	Leu Pro Glu Asn Asp Leu Ser Glu Ile Glu Asn Lys Leu Thr Phe Gln		
	245	250	255
	Asn Leu Met Glu Trp Thr Asn Pro Arg Arg Met Thr Ser Lys Tyr Val		
20	260	265	270
	Glu Val Phe Phe Pro Gln Phe Lys Ile Glu Lys Asn Tyr Glu Met Lys		
	275	280	285
	Gln Tyr Leu Arg Ala Leu Gly Leu Lys Asp Ile Phe Asp Glu Ser Lys		
25	290	295	300
	Ala Asp Leu Ser Gly Ile Ala Ser Gly Gly Arg Leu Tyr Ile Ser Arg		
	305	310	315 320
30	Met Met His Lys Ser Tyr Ile Glu Val Thr Glu Glu Gly Thr Glu Ala		
	325	330	335
	Thr Ala Ala Thr Gly Ser Asn Ile Val Glu Lys Gln Leu Pro Gln Ser		
35	340	345	350
	Thr Leu Phe Arg Ala Asp His Pro Phe Leu Phe Val Ile Arg Lys Asp		

355

360

365

Asp Ile Ile Leu Phe Ser Gly Lys Val Ser Cys Pro

370

375

380

5

<210> 3

<211> 29

<212> DNA

10 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
synthesized degenerative primer sequence

15

<400> 3

gtgaatgctg tgtacttaaa ggcaantgn

29

20

<210> 4

<211> 17

<212> DNA

<213> Artificial Sequence

25

<220>

<223> Description of Artificial Sequence:Artificially
synthesized degenerative primer sequence

<400> 4

30

aanagraang grtcngc

17

<210> 5

<211> 27

35

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
synthesized degenerative primer sequence

5

<400> 5

tatggcntcn gcngcngcng cnaaygc

27

10 <210> 6

<211> 37

<212> DNA

<213> Artificial Sequence

15 <220>

<223> Description of Artificial Sequence:Artificially
synthesized degenerative primer sequence

<400> 6

20 cgacctccag aggcaattcc agagagatca gccctgg

37

<210> 7

<211> 34

25 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
synthesized degenerative primer sequence

30

<400> 7

gtcttccaag cctacagatt tcaagtggct cctc

34

35

<210> 8

<211> 30

<212> DNA

<213> Artificial Sequence

5 <220>

<223> Description of Artificial Sequence:Artificially
synthesized antisense primer sequence

<400> 8

10 gctcagggca gtgaagatgc tcagggaaga

30

<210> 9

<211> 27

15 <212> DNA

<213> Artificial Sequence

<220>

20 <223> Description of Artificial Sequence:Artificially
synthesized antisense primer sequence

<400> 9

ctgacgtgca cagtcacctc gagcacc

27

25

<210> 10

<211> 36

<212> DNA

<213> Artificial Sequence

30

<220>

<223> Description of Artificial Sequence:Artificially
synthesized sense primer sequence

35 <400> 10

gaggtctcag aagaaggcac tgaggcaact gctgcc

36

<210> 11

<211> 16

5 <212> PRT

<213> Artificial Sequence

<220>

10 <223> Description of Artificial Sequence:Artificially
synthesized domain peptide of human megsin

<400> 11

Cys Phe Arg Glu Met Asp Asp Asn Gln Gly Asn Gly Asn Val Phe Phe
1 5 10 15

15

<210> 12

<211> 16

<212> PRT

20 <213> Artificial Sequence

<220>

25 <223> Description of Artificial Sequence:Artificially
synthesized domain peptide of human megsin

<400> 12

Cys Ser Gln Ser Gly Leu Gln Ser Gln Leu Lys Arg Val Phe Ser Asp
1 5 10 15

30

<210> 13

<211> 17

<212> PRT

<213> Artificial Sequence

35

<220>

<223> Description of Artificial Sequence:Artificially
synthesized domain peptide of human megsin

<400> 13

5 Ala Thr Gly Ser Asn Ile Val Glu Lys Gln Leu Pro Gln Ser Thr Leu
1 5 10 15

Cys

10

<210> 14

<211> 16

<212> PRT

15 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Artificially
synthesized domain peptide of human megsin

20

<400> 14

Cys Ser Asn Ile Val Glu Lys Gln Leu Pro Gln Ser Thr Leu Phe Arg
1 5 10 15

25

<210> 15

<211> 1229

<212> DNA

<213> Rattus norvegicus

30

<220>

<221> CDS

<222> (8).. (1147)

35

<300>

<310> PCT/JP98/04269

<311> 1998-09-22

<400> 15

	tttcaaa atg gcc tcc ctt gct gca gca aat gca gaa ttt ggc ttc gac	49
5	Met Ala Ser Leu Ala Ala Ala Asn Ala Glu Phe Gly Phe Asp	
	1 5 10	
	tta ttc aga gag atg gat agt agt caa gga aac gga aat gta ttc ttc	97
	Leu Phe Arg Glu Met Asp Ser Ser Gln Gly Asn Gly Asn Val Phe Phe	
10	15 20 25 30	
	tct tcc ctg agc atc ttc act gcc ctg agc cta atc cgt ttg ggt gct	145
	Ser Ser Leu Ser Ile Phe Thr Ala Leu Ser Leu Ile Arg Leu Gly Ala	
	35 40 45	
15	cga ggt gac tgt nnn cgt cag att gac aag gcc ctg cac ttt atc tcc	193
	Arg Gly Asp Cys Xaa Arg Gln Ile Asp Lys Ala Leu His Phe Ile Ser	
	50 55 60	
20	cca tca aga caa ggg aat tca tgc aac agt cag cta gga ctg caa tat	241
	Pro Ser Arg Gln Gly Asn Ser Ser Asn Ser Gln Leu Gly Leu Gln Tyr	
	65 70 75	
	caa ttg aaa aga gtt ctt gct gac ata aac tca tct cat aag gat nnn	289
25	Gln Leu Lys Arg Val Leu Ala Asp Ile Asn Ser Ser His Lys Asp Xaa	
	80 85 90	
	aaa ctc agc att gcc aat gga gtt ttt gca gag aaa gta ttt gat ttt	337
	Lys Leu Ser Ile Ala Asn Gly Val Phe Ala Glu Lys Val Phe Asp Phe	
30	95 100 105 110	
	cat aag agc tat atg gag tgt gct gaa aac tta tac aat gct aaa gtg	385
	His Lys Ser Tyr Met Glu Cys Ala Glu Asn Leu Tyr Asn Ala Lys Val	
	115 120 125	
35	gaa aga gtt gat ttt aca aat gat ata caa gaa acc aga ttt aaa att	433

	Glu Arg Val Asp Phe Thr Asn Asp Ile Gln Glu Thr Arg Phe Lys Ile	
	130 135 140	
	aat aaa tgg att gaa aat gaa aca cat ggc aaa atc aag aag gtg ttg	481
5	Asn Lys Trp Ile Glu Asn Glu Thr His Gly Lys Ile Lys Lys Val Leu	
	145 150 155	
	ggg gac agc agc ctc agc tca tca gct gtc atg gtg cta gtg aat gct	529
10	Gly Asp Ser Ser Leu Ser Ser Ser Ala Val Met Val Leu Val Asn Ala	
	160 165 170	
	gtt tac ttc aaa ggc aag tgg aaa tcg gcc ttc acc aag agt gat acc	577
15	Val Tyr Phe Lys Gly Lys Trp Lys Ser Ala Phe Thr Lys Ser Asp Thr	
	175 180 185 190	
	ctc agt tgc cat ttc agg tct ccc agc ggt cct gga aaa gca gtt aat	625
	Leu Ser Cys His Phe Arg Ser Pro Ser Gly Pro Gly Lys Ala Val Asn	
	195 200 205	
20	atg atg cat caa gaa cgg agg ttc aat ttg tct acc att cag gag cca	673
	Met Met His Gln Glu Arg Arg Phe Asn Leu Ser Thr Ile Gln Glu Pro	
	210 215 220	
	cca atg cag att ctt gag cta caa tat cat ggt ggc ata agc atg tac	721
25	Pro Met Gln Ile Leu Glu Leu Gln Tyr His Gly Gly Ile Ser Met Tyr	
	225 230 235	
	atc atg ttg ccc gag gat gac cta tcc gaa att gaa agc aag ctg agt	769
30	Ile Met Leu Pro Glu Asp Asp Leu Ser Glu Ile Glu Ser Lys Leu Ser	
	240 245 250	
	ttc cag aat cta atg gac tgg aca aat agc agg aag atg aaa tct cag	817
35	Phe Gln Asn Leu Met Asp Trp Thr Asn Ser Arg Lys Met Lys Ser Gln	
	255 260 265 270	
	tat gtg aat gtg ttt ctc ccc cag ttc aag ata gag aaa gat tat gaa	865

<213> Rattus norvegicus

<400> 16

	Met	Ala	Ser	Leu	Ala	Ala	Ala	Asn	Ala	Glu	Phe	Gly	Phe	Asp	Leu	Phe
	1				5					10					15	
5	Arg	Glu	Met	Asp	Ser	Ser	Gln	Gly	Asn	Gly	Asn	Val	Phe	Phe	Ser	Ser
				20					25					30		
	Leu	Ser	Ile	Phe	Thr	Ala	Leu	Ser	Leu	Ile	Arg	Leu	Gly	Ala	Arg	Gly
10			35						40					45		
	Asp	Cys	Xaa	Arg	Gln	Ile	Asp	Lys	Ala	Leu	His	Phe	Ile	Ser	Pro	Ser
		50					55					60				
15	Arg	Gln	Gly	Asn	Ser	Ser	Asn	Ser	Gln	Leu	Gly	Leu	Gln	Tyr	Gln	Leu
	65					70					75				80	
	Lys	Arg	Val	Leu	Ala	Asp	Ile	Asn	Ser	Ser	His	Lys	Asp	Xaa	Lys	Leu
					85					90					95	
20	Ser	Ile	Ala	Asn	Gly	Val	Phe	Ala	Glu	Lys	Val	Phe	Asp	Phe	His	Lys
				100					105					110		
	Ser	Tyr	Met	Glu	Cys	Ala	Glu	Asn	Leu	Tyr	Asn	Ala	Lys	Val	Glu	Arg
25			115					120					125			
	Val	Asp	Phe	Thr	Asn	Asp	Ile	Gln	Glu	Thr	Arg	Phe	Lys	Ile	Asn	Lys
		130					135					140				
30	Trp	Ile	Glu	Asn	Glu	Thr	His	Gly	Lys	Ile	Lys	Lys	Val	Leu	Gly	Asp
	145					150					155				160	
	Ser	Ser	Leu	Ser	Ser	Ser	Ala	Val	Met	Val	Leu	Val	Asn	Ala	Val	Tyr
					165					170					175	
35	Phe	Lys	Gly	Lys	Trp	Lys	Ser	Ala	Phe	Thr	Lys	Ser	Asp	Thr	Leu	Ser

	180		185		190
	Cys His Phe Arg Ser Pro Ser Gly Pro Gly Lys Ala Val Asn Met Met				
	195		200		205
5	His Gln Glu Arg Arg Phe Asn Leu Ser Thr Ile Gln Glu Pro Pro Met				
	210		215		220
	Gln Ile Leu Glu Leu Gln Tyr His Gly Gly Ile Ser Met Tyr Ile Met				
10	225		230		235 240
	Leu Pro Glu Asp Asp Leu Ser Glu Ile Glu Ser Lys Leu Ser Phe Gln				
		245		250	255
15	Asn Leu Met Asp Trp Thr Asn Ser Arg Lys Met Lys Ser Gln Tyr Val				
	260		265		270
	Asn Val Phe Leu Pro Gln Phe Lys Ile Glu Lys Asp Tyr Glu Met Arg				
	275		280		285
20	Ser His Leu Lys Ser Val Gly Leu Glu Asp Ile Phe Val Glu Ser Arg				
	290		295		300
	Ala Asp Leu Ser Gly Ile Ala Ser Gly Gly Arg Leu Tyr Val Ser Lys				
25	305		310		315 320
	Leu Met His Lys Ser Leu Ile Glu Val Ser Glu Glu Gly Thr Glu Ala				
		325		330	335
30	Thr Ala Ala Thr Glu Ser Asn Ile Val Glu Lys Leu Leu Pro Glu Ser				
	340		345		350
	Thr Val Phe Arg Ala Asp Arg Pro Phe Leu Phe Val Ile Arg Lys Asn				
	355		360		365
35	Gly Ile Ile Leu Phe Thr Gly Lys Val Ser Cys Pro				

370

375

380

<210> 17

5 <211> 1147

<212> DNA

<213> Mus musculus

<220>

10 <221> CDS

<222> (1).. (1104)

<300>

<310> PCT/JP98/04269

15 <311> 1998-09-22

<400> 17

ttc gac tta ttc aga gag atg gat agt agc caa gga aat gga aat gta 48

Phe Asp Leu Phe Arg Glu Met Asp Ser Ser Gln Gly Asn Gly Asn Val

20 1 5 10 15

ttc ttc tct tcc ctg agc atc ttc act gcc ctg acc cta atc cgt ctg 96

Phe Phe Ser Ser Leu Ser Ile Phe Thr Ala Leu Thr Leu Ile Arg Leu

20 25 30

25

ggt gct cga ggt gac tgt gca cgt cag att gac aag gca ctg cac ttt 144

Gly Ala Arg Gly Asp Cys Ala Arg Gln Ile Asp Lys Ala Leu His Phe

35 40 45

30 aac ata cca tca aga caa gga aac tca tct aat aat cag cca gga ctt 192

Asn Ile Pro Ser Arg Gln Gly Asn Ser Ser Asn Asn Gln Pro Gly Leu

50 55 60

cag tat caa ttg aaa aga gtt ctt gct gac ata aac tca tct cat aag 240

35 Gln Tyr Gln Leu Lys Arg Val Leu Ala Asp Ile Asn Ser Ser His Lys

65 70 75 80

	gat tat gaa ctc agc att gcc act gga gtt ttt gca gaa aaa gtc tat	288
	Asp Tyr Glu Leu Ser Ile Ala Thr Gly Val Phe Ala Glu Lys Val Tyr	
	85 90 95	
5		
	gac ttt cat aag aac tac att gag tgt gct gaa aac tta tac aat gct	336
	Asp Phe His Lys Asn Tyr Ile Glu Cys Ala Glu Asn Leu Tyr Asn Ala	
	100 105 110	
10		
	aaa gtg gaa aga gtt gac ttc aca aat gat gta caa gat acc aga ttt	384
	Lys Val Glu Arg Val Asp Phe Thr Asn Asp Val Gln Asp Thr Arg Phe	
	115 120 125	
	aaa att aat aaa tgg att gaa aat gag aca cat gga aag atc aag aag	432
15	Lys Ile Asn Lys Trp Ile Glu Asn Glu Thr His Gly Lys Ile Lys Lys	
	130 135 140	
	gtg ttg ggc gac agc agc ctc agc tcg tcg gct gtc atg gtg ctg gtg	480
	Val Leu Gly Asp Ser Ser Leu Ser Ser Ser Ala Val Met Val Leu Val	
20	145 150 155 160	
	aac gct gtt tac ttc aaa ggc aaa tgg aaa tcg gcc ttc acc aag act	528
	Asn Ala Val Tyr Phe Lys Gly Lys Trp Lys Ser Ala Phe Thr Lys Thr	
	165 170 175	
25		
	gat acc ctc agt tgc cgt ttt agg tct ccc acg tgt cct gga aaa gta	576
	Asp Thr Leu Ser Cys Arg Phe Arg Ser Pro Thr Cys Pro Gly Lys Val	
	180 185 190	
30		
	gtt aat atg atg cat caa gaa cgg cgg ttc aat ttg tct acc att cag	624
	Val Asn Met Met His Gln Glu Arg Arg Phe Asn Leu Ser Thr Ile Gln	
	195 200 205	
	cag cca cca atg cag gtt ctt gag ctc caa tat cat ggt ggc ata agc	672
35	Gln Pro Pro Met Gln Val Leu Glu Leu Gln Tyr His Gly Gly Ile Ser	
	210 215 220	

atg tac att atg ctg cct gag gat ggc cta tgt gaa att gaa agc aag 720
 Met Tyr Ile Met Leu Pro Glu Asp Gly Leu Cys Glu Ile Glu Ser Lys
 225 230 235 240

5

ctg agt ttc cag aat ctg atg gac tgg acc aat agg agg aaa atg aaa 768
 Leu Ser Phe Gln Asn Leu Met Asp Trp Thr Asn Arg Arg Lys Met Lys
 245 250 255

10

tct cag tat gtg aac gtg ttt ctc ccc cag ttc aag ata gag aag aat 816
 Ser Gln Tyr Val Asn Val Phe Leu Pro Gln Phe Lys Ile Glu Lys Asn
 260 265 270

15

tat gaa atg acg cac cac ttg aaa tcc tta ggc ttg aaa gat atc ttt 864
 Tyr Glu Met Thr His His Leu Lys Ser Leu Gly Leu Lys Asp Ile Phe
 275 280 285

20

gat gag tcc agt gca gat ctc tct gga att gcc tct gga ggt cgt ctc 912
 Asp Glu Ser Ser Ala Asp Leu Ser Gly Ile Ala Ser Gly Gly Arg Leu
 290 295 300

25

tac gta tca aag cta atg cac aag tca ttc ata gag gtc tca gag gag 960
 Tyr Val Ser Lys Leu Met His Lys Ser Phe Ile Glu Val Ser Glu Glu
 305 310 315 320

30

ggc act gaa gcc act gct gcc aca gaa aat aac att gtt gaa aag cag 1008
 Gly Thr Glu Ala Thr Ala Ala Thr Glu Asn Asn Ile Val Glu Lys Gln
 325 330 335

35

ctt cct gag tcc aca gtg ttc aga gcc gac cgc ccc ttt ctg ttt gtc 1056
 Leu Pro Glu Ser Thr Val Phe Arg Ala Asp Arg Pro Phe Leu Phe Val
 340 345 350

atc aag aag aat gac atc atc tta ttt act ggc aaa gtc tct tgt cct 1104
 Ile Lys Lys Asn Asp Ile Ile Leu Phe Thr Gly Lys Val Ser Cys Pro
 355 360 365

tgaaattcga ttgtgtttcc tatacagtaa caggcatcaa gaa

1147

5 <210> 18
 <211> 368
 <212> PRT
 <213> Mus musculus

10 <400> 18
 Phe Asp Leu Phe Arg Glu Met Asp Ser Ser Gln Gly Asn Gly Asn Val
 1 5 10 15
 Phe Phe Ser Ser Leu Ser Ile Phe Thr Ala Leu Thr Leu Ile Arg Leu
 15 20 25 30
 Gly Ala Arg Gly Asp Cys Ala Arg Gln Ile Asp Lys Ala Leu His Phe
 35 40 45
 20 Asn Ile Pro Ser Arg Gln Gly Asn Ser Ser Asn Asn Gln Pro Gly Leu
 50 55 60
 Gln Tyr Gln Leu Lys Arg Val Leu Ala Asp Ile Asn Ser Ser His Lys
 65 70 75 80
 25 Asp Tyr Glu Leu Ser Ile Ala Thr Gly Val Phe Ala Glu Lys Val Tyr
 85 90 95
 Asp Phe His Lys Asn Tyr Ile Glu Cys Ala Glu Asn Leu Tyr Asn Ala
 30 100 105 110
 Lys Val Glu Arg Val Asp Phe Thr Asn Asp Val Gln Asp Thr Arg Phe
 115 120 125
 35 Lys Ile Asn Lys Trp Ile Glu Asn Glu Thr His Gly Lys Ile Lys Lys
 130 135 140

	Val	Leu	Gly	Asp	Ser	Ser	Leu	Ser	Ser	Ser	Ala	Val	Met	Val	Leu	Val	
	145						150				155					160	
5	Asn	Ala	Val	Tyr	Phe	Lys	Gly	Lys	Trp	Lys	Ser	Ala	Phe	Thr	Lys	Thr	
						165				170					175		
	Asp	Thr	Leu	Ser	Cys	Arg	Phe	Arg	Ser	Pro	Thr	Cys	Pro	Gly	Lys	Val	
				180					185					190			
10	Val	Asn	Met	Met	His	Gln	Glu	Arg	Arg	Phe	Asn	Leu	Ser	Thr	Ile	Gln	
			195					200					205				
	Gln	Pro	Pro	Met	Gln	Val	Leu	Glu	Leu	Gln	Tyr	His	Gly	Gly	Ile	Ser	
15		210					215					220					
	Met	Tyr	Ile	Met	Leu	Pro	Glu	Asp	Gly	Leu	Cys	Glu	Ile	Glu	Ser	Lys	
	225				230					235					240		
20	Leu	Ser	Phe	Gln	Asn	Leu	Met	Asp	Trp	Thr	Asn	Arg	Arg	Lys	Met	Lys	
					245					250					255		
	Ser	Gln	Tyr	Val	Asn	Val	Phe	Leu	Pro	Gln	Phe	Lys	Ile	Glu	Lys	Asn	
				260					265					270			
25	Tyr	Glu	Met	Thr	His	His	Leu	Lys	Ser	Leu	Gly	Leu	Lys	Asp	Ile	Phe	
			275					280					285				
	Asp	Glu	Ser	Ser	Ala	Asp	Leu	Ser	Gly	Ile	Ala	Ser	Gly	Gly	Arg	Leu	
30		290					295					300					
	Tyr	Val	Ser	Lys	Leu	Met	His	Lys	Ser	Phe	Ile	Glu	Val	Ser	Glu	Glu	
	305					310					315				320		
35	Gly	Thr	Glu	Ala	Thr	Ala	Ala	Thr	Glu	Asn	Asn	Ile	Val	Glu	Lys	Gln	
					325					330					335		

Leu Pro Glu Ser Thr Val Phe Arg Ala Asp Arg Pro Phe Leu Phe Val
 340 345 350

5 Ile Lys Lys Asn Asp Ile Ile Leu Phe Thr Gly Lys Val Ser Cys Pro
 355 360 365

[Brief Description of the Drawings]

[Fig. 1]

10 Fig. 1 shows the result of Western blotting using a polyclonal antibody specific to synthetic peptide-2, which has a partial amino acid sequence of megsin protein (SEQ ID NO: 12), as the antigen. Each lane represents the following:

1: MBP

15 2: MBP-megsin protein

3: His-megsin protein

4: MBP-PAI II

5: megsin protein expressed in CHO cells

6: body fluid of the silkworm infected with a recombinant virus
 20 transfected with the megsin protein gene

7: body fluid of the silkworm infected with a recombinant virus without transfection by megsin protein gene

8: body fluid of the virus-uninfected silkworm

9: normal human serum

25 [Fig. 2]

Fig. 2 shows the result of Western blotting using a polyclonal antibody specific to synthetic peptide-3, which has a partial amino acid sequence of megsin protein (SEQ ID NO: 13), as the antigen. Each lane represents the same proteins as in Fig. 1.

30 [Fig. 3]

Fig. 3 shows the result of Western blotting using a polyclonal antibody specific to synthetic peptide-342, which has a partial amino acid sequence of megsin protein (SEQ ID NO: 15), as the antigen. Each lane represents the same proteins as in Fig. 1.

35 [Fig. 4]

Fig. 4 is a graph showing assay results of urinary megsin

proteins by ELISA. The vertical axis shows assayed value of megsin protein (creatinine ratio: AU). Urine from normal healthy person (16 samples), from patients with IgA nephropathy (24 samples), and from patients with minimal-change nephritic syndrome (24 samples) was used as specimens.

[Fig. 5]

Fig. 5 shows the assay result of Western blotting on urinary megsin protein. Each lane represents the following:

1: MBP-megsin protein

2 to 5: concentrated urine from normal healthy persons

6 and 7: concentrated urine from patients with IgA nephropathy

[Fig. 1]

1 2 3 4 5 6 7 8 9


107 —

74 —


49.3 —

36.4 —


28.5 —




[Fig. 2]




[Fig. 3]



[Fig. 4]



[Fig. 5]



[Document Name] Abstract

[Abstract]

[Problems to Be Solved] The present invention provides a method that enables diagnosis of renal diseases and determination of their severity *in vitro*.

[Means to Solve the Problems] The present invention provides a method for evaluating renal functions by measuring megsin protein in a biological specimen. Moreover, the present invention provides immunological assay reagents, as well as kits useful for the measurement of megsin protein in the biological specimen.

[Selected Drawings] None